

Product datasheet for RC224020L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Peroxiredoxin 5 (PRDX5) (NM 181651) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Peroxiredoxin 5 (PRDX5) (NM_181651) Human Tagged ORF Clone Lentiviral Particle

Symbol: PRDX5

Synonyms: ACR1; AOEB166; B166; HEL-S-55; PLP; PMP20; PRDX6; prx-V; PRXV; SBBI10

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_181651

ORF Size: 510 bp

ORF Nucleotide

Sequence:

The ORF insert of this clone is exactly the same as(RC224020).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 181651.2, NP 857634.1

 RefSeq Size:
 781 bp

 RefSeq ORF:
 513 bp

 Locus ID:
 25824

 UniProt ID:
 P30044

 Cytogenetics:
 11q13.1

Protein Families: Druggable Genome

MW: 17.4 kDa





Peroxiredoxin 5 (PRDX5) (NM_181651) Human Tagged ORF Clone Lentiviral Particle – RC224020L4V

Gene Summary:

This gene encodes a member of the peroxiredoxin family of antioxidant enzymes, which reduce hydrogen peroxide and alkyl hydroperoxides. The encoded protein interacts with peroxisome receptor 1 and plays an antioxidant protective role in different tissues under normal conditions and during inflammatory processes. The use of alternate transcription start sites is thought to result in transcript variants that use different in-frame translational start codons to generate isoforms that are targeted to the mitochondrion (isoform L) or peroxisome/cytoplasm (isoform S). Multiple related pseudogenes have been defined for this gene. [provided by RefSeq, Nov 2017]