

Product datasheet for RC224051L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

TMS1 (PYCARD) (NM_145182) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: TMS1 (PYCARD) (NM_145182) Human Tagged ORF Clone Lentiviral Particle

Symbol: PYCARD

Synonyms: ASC; CARD5; TMS; TMS-1; TMS1

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_145182

ORF Size: 528 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC224051).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 145182.2, NP 660183.1

RefSeq Size: 725 bp
RefSeq ORF: 531 bp
Locus ID: 29108
UniProt ID: Q9ULZ3
Cytogenetics: 16p11.2

Domains: PAAD_DAPIN

Protein Families: Druggable Genome





TMS1 (PYCARD) (NM_145182) Human Tagged ORF Clone Lentiviral Particle - RC224051L4V

Protein Pathways: Cytosolic DNA-sensing pathway, NOD-like receptor signaling pathway

MW: 20 kDa

Gene Summary: This gene encodes an adaptor protein that is composed of two protein-protein interaction

domains: a N-terminal PYRIN-PAAD-DAPIN domain (PYD) and a C-terminal caspase-recruitment domain (CARD). The PYD and CARD domains are members of the six-helix bundle death domain-fold superfamily that mediates assembly of large signaling complexes in the inflammatory and apoptotic signaling pathways via the activation of caspase. In normal cells, this protein is localized to the cytoplasm; however, in cells undergoing apoptosis, it forms ball-like aggregates near the nuclear periphery. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008]