

Product datasheet for RC224226L1V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Oxytocin neurophysin 1 (OXT) (NM_000915) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Oxytocin neurophysin 1 (OXT) (NM_000915) Human Tagged ORF Clone Lentiviral Particle

Symbol: Oxytocin neurophysin 1

Synonyms: OT; OT-NPI; OXT-NPI

Mammalian Cell

Selection:

None

Vector: pLenti-C-Myc-DDK (PS100064)

Tag: Myc-DDK

ACCN: NM_000915

ORF Size: 375 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC224226).

Sequence:
OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 000915.2

RefSeq Size: 512 bp
RefSeq ORF: 378 bp
Locus ID: 5020
UniProt ID: P01178
Cytogenetics: 20p13

Protein Families: Secreted Protein

MW: 12.72 kDa





Oxytocin neurophysin 1 (OXT) (NM_000915) Human Tagged ORF Clone Lentiviral Particle – RC224226L1V

Gene Summary:

This gene encodes a precursor protein that is processed to produce oxytocin and neurophysin I. Oxytocin is a posterior pituitary hormone which is synthesized as an inactive precursor in the hypothalamus along with its carrier protein neurophysin I. Together with neurophysin, it is packaged into neurosecretory vesicles and transported axonally to the nerve endings in the neurohypophysis, where it is either stored or secreted into the bloodstream. The precursor seems to be activated while it is being transported along the axon to the posterior pituitary. This hormone contracts smooth muscle during parturition and lactation. It is also involved in cognition, tolerance, adaptation and complex sexual and maternal behaviour, as well as in the regulation of water excretion and cardiovascular functions. [provided by RefSeq, Dec 2013]