

Product datasheet for RC224322L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200
Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

WDR19 (NM_025132) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Symbol: WDR19

Synonyms: ATD5; CED4; DYF-2; FAP66; IFT144; NPHP13; ORF26; Oseg6; PWDMP; SRTD5

Mammalian Cell Puromycin

Selection:

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_025132

ORF Size: 4026 bp

ORF Nucleotide Sequence: The ORF insert of this clone is exactly the same as(RC224322).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM_025132.3</u>

RefSeq Size: 4534 bp

RefSeq ORF: 4029 bp

Locus ID: 57728

UniProt ID: Q8NEZ3

Cytogenetics: 4pl4





WDR19 (NM_025132) Human Tagged ORF Clone Lentiviral Particle | RC224322L4V

Domains: WD40

MW: 151.4 kDa

Gene Summary:

The protein encoded by this gene is a member of the WD (tryptophan-aspartic acid) repeat family, which is a large family of structurally-related proteins known to participate in a wide range of cellular processes. Each WD repeat typically contains about 40 amino acids that are usually bracketed by glycine-histidine and tryptophan-aspartic acid (WD) dipeptides. This protein contains six WD repeats, three transmembrane domains, and a clathrin heavy-chain repeat. Mutations in this gene have been described in individuals with a wide range of disorders affecting function of the cilium. These disorders are known as ciliopathies, and include Jeune syndrome, Sensenbrenner syndromes, Senior-Loken syndrome, combined or isolated nephronophthisis (NPHP), and retinitis pigmentosa (RP). Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Dec 2015]