

## Product datasheet for RC224361L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## HNMT (NM\_001024075) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** HNMT (NM\_001024075) Human Tagged ORF Clone Lentiviral Particle

Symbol: HNMT

Synonyms: HMT; HNMT-S1; HNMT-S2; MRT51

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK

**ACCN:** NM\_001024075

ORF Size: 378 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC224361).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 001024075.1

 RefSeq Size:
 907 bp

 RefSeq ORF:
 381 bp

 Locus ID:
 3176

 UniProt ID:
 P50135

 Cytogenetics:
 2q22.1

**Protein Families:** Druggable Genome

**Protein Pathways:** Histidine metabolism





ORIGENE

**MW:** 14 kDa

**Gene Summary:** 

In mammals, histamine is metabolized by two major pathways: N(tau)-methylation via histamine N-methyltransferase and oxidative deamination via diamine oxidase. This gene encodes the first enzyme which is found in the cytosol and uses S-adenosyl-L-methionine as the methyl donor. In the mammalian brain, the neurotransmitter activity of histamine is controlled by N(tau)-methylation as diamine oxidase is not found in the central nervous system. A common genetic polymorphism affects the activity levels of this gene product in red blood cells. Multiple alternatively spliced transcript variants that encode different proteins have been found for this gene. [provided by RefSeq, Jul 2008]