

### Product datasheet for RC224383L4

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Killer cell immunoglobulin like receptor 3DL3 (KIR3DL3) (NM\_153443) Human Tagged Lenti ORF Clone

### **Product data:**

**Product Type:** Expression Plasmids

Product Name: Killer cell immunoglobulin like receptor 3DL3 (KIR3DL3) (NM\_153443) Human Tagged Lenti

**ORF Clone** 

Tag: mGFP

Symbol: Killer cell immunoglobulin like receptor 3DL3

Synonyms: CD158Z; KIR3DL7; KIR44; KIRC1

**Mammalian Cell** 

Selection:

Puromycin

**Vector:** pLenti-C-mGFP-P2A-Puro (PS100093)

E. coli Selection: Chloramphenicol (34 ug/mL)


**ORF Nucleotide** 


The ORF insert of this clone is exactly the same as(RC224383).

Sequence:

**Restriction Sites:** Sgfl-Rsrll

**Cloning Scheme:** 





 $<sup>\</sup>ensuremath{^*}$  The last codon before the Stop codon of the ORF.

**ACCN:** NM 153443





Killer cell immunoglobulin like receptor 3DL3 (KIR3DL3) (NM\_153443) Human Tagged Lenti ORF Clone - RC224383L4

ORF Size: 1230 bp

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

**RefSeq:** <u>NM 153443.2</u>, <u>NP 703144.2</u>

RefSeq Size: 1691 bp
RefSeq ORF: 1233 bp
Locus ID: 115653
Cytogenetics: 19q13.42

**Protein Families:** Transmembrane

**Protein Pathways:** Antigen processing and presentation


**MW:** 44.7 kDa

**Gene Summary:** Killer cell immunoglobulin-like receptors (KIRs) are transmembrane glycoproteins expressed

by natural killer cells and subsets of T cells. The KIR genes are polymorphic and highly homologous and they are found in a cluster on chromosome 19q13.4 within the 1 Mb leukocyte receptor complex (LRC). The gene content of the KIR gene cluster varies among haplotypes, although several "framework" genes are found in all haplotypes (KIR3DL3, KIR3DP1, KIR3DL4, KIR3DL2). The KIR proteins are classified by the number of extracellular immunoglobulin domains (2D or 3D) and by whether they have a long (L) or short (S) cytoplasmic domain. KIR proteins with the long cytoplasmic domain transduce inhibitory signals upon ligand binding via an immune tyrosine-based inhibitory motif (ITIM), while KIR proteins with the short cytoplasmic domain lack the ITIM motif and instead associate with the TYRO protein tyrosine kinase binding protein to transduce activating signals. The ligands for several KIR proteins are subsets of HLA class I molecules; thus, KIR proteins are thought to play an important role in regulation of the immune response. This gene is one of the "framework" loci that is present on all haplotypes. [provided by RefSeq, Jul 2008]



## **Product images:**



Circular map for RC224383L4