

Product datasheet for RC224398L4

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

ErbB 3 (ERBB3) (NM_001005915) Human Tagged Lenti ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: ErbB 3 (ERBB3) (NM_001005915) Human Tagged Lenti ORF Clone

Tag: mGFP Symbol: ErbB 3

Synonyms: c-erbB-3; c-erbB3; ErbB-3; erbB3-5; FERLK; HER3; LCCS2; MDA-BF-1; p45-sErbB3; p85-sErbB3;

p180-ErbB3

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

E. coli Selection: Chloramphenicol (34 ug/mL)

ORF Nucleotide


The ORF insert of this clone is exactly the same as(RC224398).

Sequence:

Restriction Sites: Sgfl-Mlul

Cloning Scheme:





^{*} The last codon before the Stop codon of the ORF.

ACCN: NM 001005915

ORF Size: 549 bp



ErbB 3 (ERBB3) (NM_001005915) Human Tagged Lenti ORF Clone - RC224398L4

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001005915.1</u>, <u>NP 001005915.1</u>

12q13.2

 RefSeq Size:
 1050 bp

 RefSeq ORF:
 552 bp

 Locus ID:
 2065

 UniProt ID:
 P21860

Cytogenetics:

Protein Families: Adult stem cells, Druggable Genome, Protein Kinase, Secreted Protein, Stem cell -

Pluripotency, Transmembrane

Protein Pathways: Calcium signaling pathway, Endocytosis, ErbB signaling pathway

MW: 20.17 kDa

Gene Summary: This gene encodes a member of the epidermal growth factor receptor (EGFR) family of


receptor tyrosine kinases. This membrane-bound protein has a neuregulin binding domain but not an active kinase domain. It therefore can bind this ligand but not convey the signal into the cell through protein phosphorylation. However, it does form heterodimers with other EGF receptor family members which do have kinase activity. Heterodimerization leads to the activation of pathways which lead to cell proliferation or differentiation. Amplification of this gene and/or overexpression of its protein have been reported in numerous cancers, including prostate, bladder, and breast tumors. Alternate transcriptional splice variants encoding

different isoforms have been characterized. One isoform lacks the intermembrane region and is secreted outside the cell. This form acts to modulate the activity of the membrane-bound form. Additional splice variants have also been reported, but they have not been thoroughly

characterized. [provided by RefSeq, Jul 2008]



Product images:



Circular map for RC224398L4