

# Product datasheet for RC224432L3V

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## DKK2 (NM\_014421) Human Tagged ORF Clone Lentiviral Particle

#### **Product data:**

Product Type: Lentiviral Particles

Product Name: DKK2 (NM 014421) Human Tagged ORF Clone Lentiviral Particle

Symbol: DKK2
Synonyms: DKK-2

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

 Tag:
 Myc-DDK

 ACCN:
 NM\_014421

ORF Size: 777 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC224432).

Sequence:
OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 014421.2

 RefSeq Size:
 3659 bp

 RefSeq ORF:
 780 bp

 Locus ID:
 27123

 UniProt ID:
 Q9UBU2

 Cytogenetics:
 4q25

**Protein Families:** Adult stem cells, Cancer stem cells, Druggable Genome, ES Cell Differentiation/IPS, Secreted

Protein, Stem cell relevant signaling - Wnt Signaling pathway





### DKK2 (NM\_014421) Human Tagged ORF Clone Lentiviral Particle - RC224432L3V

**Protein Pathways:** Wnt signaling pathway

**MW:** 28.45 kDa

**Gene Summary:** This gene encodes a protein that is a member of the dickkopf family. The secreted protein

contains two cysteine rich regions and is involved in embryonic development through its interactions with the Wnt signaling pathway. It can act as either an agonist or antagonist of Wnt/beta-catenin signaling, depending on the cellular context and the presence of the cofactor kremen 2. Activity of this protein is also modulated by binding to the Wnt co-receptor

LDL-receptor related protein 6 (LRP6). [provided by RefSeq, Jul 2008]