

Product datasheet for RC224462L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200
Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

SORCS3 (NM_014978) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Symbol: SORCS3

Synonyms: SORCS

Mammalian Cell Puromycin

Selection:

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK

ACCN: NM_014978

ORF Size: 3666 bp

ORF Nucleotide Sequence: The ORF insert of this clone is exactly the same as(RC224462).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM_014978.1</u>

RefSeq Size: 5757 bp

RefSeq ORF: 3669 bp

Locus ID: 22986

UniProt ID: Q9UPU3

Cytogenetics: 10q25.1



SORCS3 (NM_014978) Human Tagged ORF Clone Lentiviral Particle | RC224462L3V

Protein Families: Druggable Genome, Transmembrane

MW: 135.8 kDa

Gene Summary:

This gene encodes a type-I receptor transmembrane protein that is a member of the vacuolar protein sorting 10 receptor family. Proteins of this family are defined by a vacuolar protein sorting 10 domain at the N-terminus. The N-terminal segment of this domain has a consensus motif for proprotein convertase processing, and the C-terminal segment of this domain is characterized by ten conserved cysteine residues. The vacuolar protein sorting 10 domain is followed by a leucine-rich segment, a transmembrane domain, and a short C-terminal cytoplasmic domain that interacts with adaptor molecules. The transcript is expressed at high levels in the brain, and candidate gene studies suggest that genetic variation in this gene is associated with Alzheimer's disease. Consistent with this observation, knockdown of the gene in cell culture results in an increase in amyloid precursor protein processing. [provided by RefSeq, Dec 2014]