

## Product datasheet for RC224481L1V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **OGT (NM\_181673) Human Tagged ORF Clone Lentiviral Particle**

**Product data:** 

Product Type: Lentiviral Particles

**Product Name:** OGT (NM\_181673) Human Tagged ORF Clone Lentiviral Particle

Symbol: OG1

Synonyms: HINCUT-1; HRNT1; MRX106; O-GLCNAC; OGT1

Mammalian Cell

Selection:

None

**Vector:** pLenti-C-Myc-DDK (PS100064)

Tag: Myc-DDK
ACCN: NM 181673

ORF Size: 3108 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC224481).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 181673.1

RefSeq Size: 5445 bp
RefSeq ORF: 3111 bp
Locus ID: 8473
UniProt ID: 015294
Cytogenetics: Xq13.1

**Protein Families:** Druggable Genome

**Protein Pathways:** Metabolic pathways, O-Glycan biosynthesis





## OGT (NM\_181673) Human Tagged ORF Clone Lentiviral Particle - RC224481L1V

**MW:** 115.5 kDa

Gene Summary: This

This gene encodes a glycosyltransferase that catalyzes the addition of a single N-acetylglucosamine in O-glycosidic linkage to serine or threonine residues. Since both phosphorylation and glycosylation compete for similar serine or threonine residues, the two processes may compete for sites, or they may alter the substrate specificity of nearby sites by steric or electrostatic effects. The protein contains multiple tetratricopeptide repeats that are required for optimal recognition of substrates. Alternatively spliced transcript variants encoding distinct isoforms have been found for this gene. [provided by RefSeq, Oct 2009]