

Product datasheet for RC224529L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Retinoic Acid Receptor beta (RARB) (NM_000965) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Retinoic Acid Receptor beta (RARB) (NM_000965) Human Tagged ORF Clone Lentiviral Particle

Symbol: Retinoic Acid Receptor beta

Synonyms: HAP; MCOPS12; NR1B2; RARbeta1; RRB2

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

HOLI, zf-C4

Tag: mGFP

ACCN: NM_000965

ORF Size: 1344 bp

ORF Nucleotide

Sequence:

Domains:

The ORF insert of this clone is exactly the same as(RC224529).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 000965.2</u>

 RefSeq Size:
 3222 bp

 RefSeq ORF:
 1347 bp

 Locus ID:
 5915

 UniProt ID:
 P10826

 Cytogenetics:
 3p24.2





Retinoic Acid Receptor beta (RARB) (NM_000965) Human Tagged ORF Clone Lentiviral Particle – RC224529L4V

Protein Families: Druggable Genome, Nuclear Hormone Receptor, Transcription Factors

Protein Pathways: Non-small cell lung cancer, Pathways in cancer, Small cell lung cancer

MW: 50.3 kDa

Gene Summary: This gene encodes retinoic acid receptor beta, a member of the thyroid-steroid hormone

receptor superfamily of nuclear transcriptional regulators. This receptor localizes to the cytoplasm and to subnuclear compartments. It binds retinoic acid, the biologically active form of vitamin A which mediates cellular signalling in embryonic morphogenesis, cell growth and differentiation. It is thought that this protein limits growth of many cell types by regulating gene expression. The gene was first identified in a hepatocellular carcinoma where it flanks a hepatitis B virus integration site. Alternate promoter usage and differential splicing result in

multiple transcript variants. [provided by RefSeq, Mar 2014]