

Product datasheet for RC224537L2V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Apc11 (ANAPC11) (NM 016476) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Apc11 (ANAPC11) (NM_016476) Human Tagged ORF Clone Lentiviral Particle

Symbol: Apc11

Synonyms: APC11; Apc11p; HSPC214

Mammalian Cell

Selection:

None

Vector: pLenti-C-mGFP (PS100071)

Tag: mGFP

ACCN: NM_016476

ORF Size: 252 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC224537).

Sequence:

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 016476.10

RefSeq Size: 892 bp
RefSeq ORF: 255 bp
Locus ID: 51529
UniProt ID: Q9NYG5
Cytogenetics: 17q25.3

Protein Families: Druggable Genome





Apc11 (ANAPC11) (NM_016476) Human Tagged ORF Clone Lentiviral Particle - RC224537L2V

Protein Pathways: Cell cycle, Oocyte meiosis, Progesterone-mediated oocyte maturation, Ubiquitin mediated

proteolysis

MW: 9.7 kDa

Gene Summary: Together with the cullin protein ANAPC2, constitutes the catalytic component of the anaphase

promoting complex/cyclosome (APC/C), a cell cycle-regulated E3 ubiquitin ligase that controls progression through mitosis and the G1 phase of the cell cycle. The APC/C complex acts by mediating ubiquitination and subsequent degradation of target proteins: it mainly mediates the formation of 'Lys-11'-linked polyubiquitin chains and, to a lower extent, the formation of 'Lys-48'- and 'Lys-63'-linked polyubiquitin chains. May recruit the E2 ubiquitin-conjugating

enzymes to the complex.[UniProtKB/Swiss-Prot Function]