

## Product datasheet for RC224556L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Fibrinogen alpha chain (FGA) (NM\_021871) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** Fibrinogen alpha chain (FGA) (NM\_021871) Human Tagged ORF Clone Lentiviral Particle

Symbol: Fibrinogen alpha chain

Synonyms: Fib2

Mammalian Cell

Puromycin

Selection:

Vector:

pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK
ACCN: NM\_021871

ORF Size: 1932 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC224556).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 021871.2

 RefSeq Size:
 2210 bp

 RefSeq ORF:
 1935 bp

 Locus ID:
 2243

 UniProt ID:
 P02671

 Cytogenetics:
 4q31.3

**Protein Families:** Druggable Genome, Secreted Protein

**Protein Pathways:** Complement and coagulation cascades





## Fibrinogen alpha chain (FGA) (NM\_021871) Human Tagged ORF Clone Lentiviral Particle – RC224556L3V

**MW:** 67.6 kDa

**Gene Summary:** 

This gene encodes the alpha subunit of the coagulation factor fibrinogen, which is a component of the blood clot. Following vascular injury, the encoded preproprotein is proteolytically processed by thrombin during the conversion of fibrinogen to fibrin. Mutations in this gene lead to several disorders, including dysfibrinogenemia, hypofibrinogenemia, afibrinogenemia and renal amyloidosis. Alternative splicing results in multiple transcript variants, at least one of which encodes an isoform that undergoes proteolytic processing. [provided by RefSeq, Jan 2016]