

Product datasheet for RC224758L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CD11d (ITGAD) (NM 005353) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: CD11d (ITGAD) (NM_005353) Human Tagged ORF Clone Lentiviral Particle

Symbol:

ADB2: CD11D Synonyms:

Mammalian Cell

Puromycin

Selection:

Vector:

pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK

NM 005353 ACCN:

ORF Size: 3483 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC224758).

Sequence:

Cytogenetics:

The molecular sequence of this clone aligns with the gene accession number as a point of OTI Disclaimer:

> reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 005353.2, NP 005344.2

RefSeq Size: 3912 bp RefSeq ORF: 3486 bp Locus ID: 3681 **UniProt ID:** Q13349

16p11.2 **Protein Families:** Druggable Genome, Transmembrane

Protein Pathways: Regulation of actin cytoskeleton





ORIGENE

MW: 126.76 kDa

Gene Summary: This gene belongs to the beta-2 integrin family of membrane glycoproteins, which are are

composed of non-covalently linked alpha and beta subunits to form a heterodimer. It encodes the alpha subunit of the cell surface heterodimers and is involved in the activation and adhesion functions of leukocytes. The gene is located about 11kb downstream of the integrin subunit alpha X gene, another member of the integrin family. It is expressed in the tissue and circulating myeloid leukocytes. Alternative splicing results in multiple transcript

variants. [provided by RefSeq, Dec 2015]