


## Product datasheet for RC224760

### SETD2 (NM\_014159) Human Tagged ORF Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	SETD2 (NM_014159) Human Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	SETD2
Synonyms:	HBP231; HIF-1; HIP-1; HSPC069; HYPB; KMT3A; LLS; p231HBP; SET2
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Chromatograms:	<a href="https://cdn.origene.com/chromatograms/mk8036_h08.zip">https://cdn.origene.com/chromatograms/mk8036_h08.zip</a>
Restriction Sites:	Sgfl-MluI
Cloning Scheme:	

Cloning sites used for ORF Shuttling:



\* The last codon before the Stop codon of the ORF


ACCN:	NM_014159
ORF Size:	7692 bp



[View online >](#)

<b>OTI Disclaimer:</b>	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
<b>OTI Annotation:</b>	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>RefSeq:</b>	<a href="#">NM_014159.7</a>
<b>RefSeq Size:</b>	8452 bp
<b>RefSeq ORF:</b>	7695 bp
<b>Locus ID:</b>	29072
<b>UniProt ID:</b>	<a href="#">Q9BYW2</a>
<b>Cytogenetics:</b>	3p21.31
<b>Domains:</b>	WW, SET, PostSET, AWS
<b>Protein Families:</b>	Druggable Genome
<b>Protein Pathways:</b>	Lysine degradation
<b>MW:</b>	287.6 kDa
<b>Gene Summary:</b>	Huntington's disease (HD), a neurodegenerative disorder characterized by loss of striatal neurons, is caused by an expansion of a polyglutamine tract in the HD protein huntingtin. This gene encodes a protein belonging to a class of huntingtin interacting proteins characterized by WW motifs. This protein is a histone methyltransferase that is specific for lysine-36 of histone H3, and methylation of this residue is associated with active chromatin. This protein also contains a novel transcriptional activation domain and has been found associated with hyperphosphorylated RNA polymerase II. [provided by RefSeq, Aug 2008]

Product images:



Circular map for RC224760