

Product datasheet for **RC224791L2V**

GDF10 (NM_004962) Human Tagged ORF Clone Lentiviral Particle

Product data:

| | |
|---------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Product Type: | Lentiviral Particles |
| Product Name: | GDF10 (NM_004962) Human Tagged ORF Clone Lentiviral Particle |
| Symbol: | GDF10 |
| Synonyms: | BIP; BMP-3b; BMP3B |
| Mammalian Cell Selection: | None |
| Vector: | pLenti-C-mGFP (PS100071) |
| Tag: | mGFP |
| ACCN: | NM_004962 |
| ORF Size: | 1434 bp |
| ORF Nucleotide Sequence: | The ORF insert of this clone is exactly the same as(RC224791). |
| OTI Disclaimer: | The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info |
| OTI Annotation: | This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene. |
| RefSeq: | NM_004962.2 |
| RefSeq Size: | 2674 bp |
| RefSeq ORF: | 1437 bp |
| Locus ID: | 2662 |
| UniProt ID: | P55107 |
| Cytogenetics: | 10q11.22 |
| Protein Families: | Druggable Genome, Secreted Protein |
| MW: | 52.9 kDa |



[View online »](#)

Gene Summary:

This gene encodes a secreted ligand of the TGF-beta (transforming growth factor-beta) superfamily of proteins. Ligands of this family bind various TGF-beta receptors leading to recruitment and activation of SMAD family transcription factors that regulate gene expression. The encoded preproprotein is proteolytically processed to generate each subunit of the disulfide-linked homodimer. This promotes neural repair after stroke. Additionally, this protein may act as a tumor suppressor and reduced expression of this gene is associated with oral cancer. [provided by RefSeq, Jul 2016]