

Product datasheet for RC224805L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Prostaglandin F2 alpha Receptor (PTGFR) (NM_000959) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Prostaglandin F2 alpha Receptor (PTGFR) (NM_000959) Human Tagged ORF Clone Lentiviral

Particle

Symbol: Prostaglandin F2 alpha Receptor

Synonyms: FP

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_000959

ORF Size: 1077 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC224805).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 000959.2</u>

 RefSeq Size:
 3466 bp

 RefSeq ORF:
 1080 bp

 Locus ID:
 5737

 UniProt ID:
 P43088

 Cytogenetics:
 1p31.1

Domains: 7tm_1





Prostaglandin F2 alpha Receptor (PTGFR) (NM_000959) Human Tagged ORF Clone Lentiviral Particle - RC224805L4V

Protein Families: Druggable Genome, GPCR, Transmembrane

Protein Pathways: Calcium signaling pathway, Neuroactive ligand-receptor interaction

MW: 40.05 kDa

Gene Summary: The protein encoded by this gene is member of the G-protein coupled receptor family. This

protein is a receptor for prostaglandin F2-alpha (PGF2-alpha), which is known to be a potent luteolytic agent, and may also be involved in modulating intraocular pressure and smooth muscle contraction in uterus. Knockout studies in mice suggest that the interaction of PGF2-alpha with this receptor may initiate parturition in ovarian luteal cells and thus induce

luteolysis. Two transcript variants encoding different isoforms have been found for this gene.

[provided by RefSeq, Jul 2008]