

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Product datasheet for RC224831L3V

## AASS (NM\_005763) Human Tagged ORF Clone Lentiviral Particle

## **Product data:**

Product Type:	Lentiviral Particles
Product Name:	AASS (NM_005763) Human Tagged ORF Clone Lentiviral Particle
Symbol:	AASS
Synonyms:	LKR/SDH; LKRSDH; LORSDH
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_005763
ORF Size:	2778 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC224831).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 005763.2</u>
RefSeq Size:	3233 bp
RefSeq ORF:	2781 bp
Locus ID:	10157
UniProt ID:	<u>Q9UDR5</u>
Cytogenetics:	7q31.32
Domains:	Saccharop_dh, AlaDh_PNT_C, AlaDh_PNT_N
Protein Families:	Druggable Genome



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Serigene AASS (NM_005763) Human Tagged ORF Clone Lentiviral Particle – RC224831L3V	
Protein Pathway	<b>/s:</b> Lysine biosynthesis, Lysine degradation, Metabolic pathways
MW:	102 kDa
Gene Summary:	This gene encodes a bifunctional enzyme that catalyzes the first two steps in the mammalian lysine degradation pathway. The N-terminal and the C-terminal portions of this enzyme contain lysine-ketoglutarate reductase and saccharopine dehydrogenase activity, respectively, resulting in the conversion of lysine to alpha-aminoadipic semialdehyde. Mutations in this gene are associated with familial hyperlysinemia. [provided by RefSeq, Jul 2008]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US