

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for RC224839L1V

FAM62B (ESYT2) (NM_020728) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	FAM62B (ESYT2) (NM_020728) Human Tagged ORF Clone Lentiviral Particle
Symbol:	FAM62B
Synonyms:	CHR2SYT; E-Syt2; FAM62B
Mammalian Cell Selection:	None
Vector:	pLenti-C-Myc-DDK (PS100064)
Tag:	Myc-DDK
ACCN:	NM_020728
ORF Size:	2679 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC224839).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 020728.1, NP 065779.1</u>
RefSeq Size:	5956 bp
RefSeq ORF:	2538 bp
Locus ID:	57488
UniProt ID:	<u>A0FGR8</u>
Cytogenetics:	7q36.3
Protein Families:	Transmembrane
MW:	98.7 kDa



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



FAM62B (ESYT2) (NM_020728) Human Tagged ORF Clone Lentiviral Particle – RC224839L1V

Gene Summary:Tethers the endoplasmic reticulum to the cell membrane and promotes the formation of
appositions between the endoplasmic reticulum and the cell membrane. Binds
glycerophospholipids in a barrel-like domain and may play a role in cellular lipid transport.
Plays a role in FGF signaling via its role in the rapid internalization of FGFR1 that has been
activated by FGF1 binding; this occurs most likely via the AP-2 complex.[UniProtKB/Swiss-Prot
Function]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US