

Product datasheet for RC224861L1V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CTBP2 (NM_001329) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: CTBP2 (NM_001329) Human Tagged ORF Clone Lentiviral Particle

Symbol: CTBP2

Mammalian Cell

Selection:

None

Vector: pLenti-C-Myc-DDK (PS100064)

Tag: Myc-DDK

ACCN: NM_001329

ORF Size: 1335 bp

ORF Nucleotide

OTI Disclaimer:

The ORF insert of this clone is exactly the same as(RC224861).

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 001329.1</u>

RefSeq Size: 2368 bp
RefSeq ORF: 1338 bp
Locus ID: 1488
UniProt ID: P56545

Cytogenetics: 10q26.13

Domains: 2-Hacid_DH, 2-Hacid_DH_C

Protein Families: Stem cell - Pluripotency, Stem cell relevant signaling - Wnt Signaling pathway





CTBP2 (NM_001329) Human Tagged ORF Clone Lentiviral Particle - RC224861L1V

Protein Pathways: Chronic myeloid leukemia, Notch signaling pathway, Pathways in cancer, Wnt signaling

pathway

MW: 48.8 kDa

Gene Summary: This gene produces alternative transcripts encoding two distinct proteins. One protein is a

transcriptional repressor, while the other isoform is a major component of specialized synapses known as synaptic ribbons. Both proteins contain a NAD+ binding domain similar to NAD+-dependent 2-hydroxyacid dehydrogenases. A portion of the 3' untranslated region was

used to map this gene to chromosome 21q21.3; however, it was noted that similar loci elsewhere in the genome are likely. Blast analysis shows that this gene is present on chromosome 10. Several transcript variants encoding two different isoforms have been

found for this gene. [provided by RefSeq, Feb 2014]