

Product datasheet for RC224959L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Phosphatidic acid phosphatase type 2B (PLPP3) (NM_177414) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Phosphatidic acid phosphatase type 2B (PLPP3) (NM_177414) Human Tagged ORF Clone

Lentiviral Particle

Symbol: Phosphatidic acid phosphatase type 2B

Synonyms: Dri42; LPP3; MGC15306; PAP-2b; PAP2-beta; VCIP

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag:Myc-DDKACCN:NM_177414

ORF Size: 933 bp

ORF Nucleotide

Sequence:

The ORF insert of this clone is exactly the same as(RC224959).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 177414.1, NP 803133.1</u>

RefSeq Size: 3243 bp
RefSeq ORF: 935 bp
Locus ID: 8613
Cytogenetics: 1p32.2

Protein Families: Druggable Genome, Transmembrane





Phosphatidic acid phosphatase type 2B (PLPP3) (NM_177414) Human Tagged ORF Clone Lentiviral Particle – RC224959L3V

Protein Pathways: Ether lipid metabolism, Fc gamma R-mediated phagocytosis, Glycerolipid metabolism,

Glycerophospholipid metabolism, Metabolic pathways, Sphingolipid metabolism

MW: 34.9 kDa

Gene Summary: The protein encoded by this gene is a member of the phosphatidic acid phosphatase (PAP)

family. PAPs convert phosphatidic acid to diacylglycerol, and function in de novo synthesis of glycerolipids as well as in receptor-activated signal transduction mediated by phospholipase D. This protein is a membrane glycoprotein localized at the cell plasma membrane. It has been shown to actively hydrolyze extracellular lysophosphatidic acid and short-chain

phosphatidic acid. The expression of this gene is found to be enhanced by epidermal growth

factor in Hela cells. [provided by RefSeq, Mar 2010]