

## Product datasheet for RC224975L1V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Legumain (LGMN) (NM\_005606) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

Product Name: Legumain (LGMN) (NM 005606) Human Tagged ORF Clone Lentiviral Particle

Symbol: Legumain

**Synonyms:** AEP; LGMN1; PRSC1

Mammalian Cell

Selection:

None

**Vector:** pLenti-C-Myc-DDK (PS100064)

Tag: Myc-DDK
ACCN: NM\_005606

ORF Size: 1299 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC224975).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 005606.5</u>

 RefSeq Size:
 2073 bp

 RefSeq ORF:
 1302 bp

 Locus ID:
 5641

 UniProt ID:
 Q99538

 Cytogenetics:
 14q32.12

**Domains:** Peptidase\_C13

**Protein Families:** Druggable Genome, Protease





**Protein Pathways:** Antigen processing and presentation, Lysosome

**MW:** 49.4 kDa

**Gene Summary:** This gene encodes a cysteine protease that has a strict specificity for hydrolysis of

asparaginyl bonds. This enzyme may be involved in the processing of bacterial peptides and endogenous proteins for MHC class II presentation in the lysosomal/endosomal systems. Enzyme activation is triggered by acidic pH and appears to be autocatalytic. Protein expression occurs after monocytes differentiate into dendritic cells. A fully mature, active enzyme is produced following lipopolysaccharide expression in mature dendritic cells. Overexpression of this gene may be associated with the majority of solid tumor types. This gene has a pseudogene on chromosome 13. Several alternatively spliced transcript variants have been described, but the biological validity of only two has been determined. These two

variants encode the same isoform. [provided by RefSeq, Jul 2008]