

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for RC225183

C9orf95 (NMRK1) (NM_001127603) Human Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	C9orf95 (NMRK1) (NM_001127603) Human Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	C9orf95
Synonyms:	bA235O14.2; C9orf95; NRK1
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
ORF Nucleotide Sequence:	<pre>>RC225183 representing NM_001127603 Red=Cloning site Blue=ORF Green=Tags(s)</pre>
	TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC GCC <mark>GCGATCGC</mark> C
	ATGAAAACATTTATCATTGGAATCAGTGGTGTGACAAACAGTGGCAAAACAACACCACGGCTAAGAATTTGC AGAAACACCTCCCAAATTGCAGTGTCATATCTCAGGATGATTTCTTCAAGCCAGAGTCTGAGATAGAGAC AGATAAAAATGGATTTTTGCAGTACGATGTGCTTGAAGCACTTAACATGGAAAAATGATGTCAGCCATT TCCTGCTGGATGGAAAGCGCAAGACACTCTGTGGTATCAACAGACCAGGAAAGTGCTGAGGAAATTCCCA TTTTAATCATCGAAGGTTTTCTTCTTTTTAATTATAATACAAGGGTCTATCAGCCTCCAGACTCTCCGGG ATACTTTGATGGCCATGTGTGGCCCATGTATCTAAAGTACAGACAAGAAATGCAGGACATCACATGGGAA GTTGTGTACCTGGATGGAACAAAATCTGAAGAGGACCTCTTTTGCAAGTATATGAAGATCTAATACAAG AACTAGCAAAGCAA
	ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT ACAAGGATGACGACGATAAG GTTTAA
Protein Sequence:	>RC225183 representing NM_001127603 <mark>Red</mark> =Cloning site Green=Tags(s)
	MKTFIIGISGVTNSGKTTLAKNLQKHLPNCSVISQDDFFKPESEIETDKNGFLQYDVLEALNMEKMMSAI SCWMESARHSVVSTDQESAEEIPILIIEGFLLFNYNTRVYQPPDSPGYFDGHVWPMYLKYRQEMQDITWE VVYLDGTKSEEDLFLQVYEDLIQELAKQKCLQVTA
	TRTRPLEQKLISEEDLAANDILDYKDDDDKV
Chromatograms:	https://cdn.origene.com/chromatograms/ja1456_e04.zip




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

C9orf95 (NMRK1) (NM_001127603) Human Tagged ORF Clone – RC225183

Restriction Sites:

Cloning Scheme:




	* The last codon before the Stop codon of the ORF
ACCN:	NM_001127603
ORF Size:	525 bp
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	 Centrifuge at 5,000xg for 5min. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. Close the tube and incubate for 10 minutes at room temperature. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM 001127603.2</u>
RefSeq ORF:	528 bp
Locus ID:	54981
UniProt ID:	<u>Q9NWW6</u>
Cytogenetics:	9q21.13


This product is to be used for laboratory only. Not for diagnostic or therapeutic use. <u>©2023 OriGene Techno</u>logies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

ORIGENE C9orf	95 (NMRK1) (NM_001127603) Human Tagged ORF Clone – RC225183
Protein Pathways:	Nicotinate and nicotinamide metabolism
MW:	20 kDa
Gene Summary:	Nicotinamide adenine dinucleotide (NAD+) is essential for life in all organisms, both as a coenzyme for oxidoreductases and as a source of ADP-ribosyl groups used in various reactions. Nicotinic acid and nicotinamide, collectively known as niacin, are the vitamin precursors of NAD+. Nicotinamide riboside kinases, such as NRK1, function to synthesize NAD+ through nicotinamide mononucleotide using nicotinamide riboside as the precursor (Bieganowski and Brenner, 2004 [PubMed 15137942]).[supplied by OMIM, Mar 2008]

Product images:



Circular map for RC225183



Western blot validation of overexpression lysate (Cat# [LY426819]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with RC225183 using transfection reagent MegaTran 2.0 (Cat# [TT210002]).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US