

Product datasheet for RC225257

NACA (NM_001113202) Human Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	NACA (NM_001113202) Human Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	NACA
Synonyms:	HSD48; NAC-alpha; NACA1; skNAC
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
ORF Nucleotide Sequence:	>RC225257 representing NM_001113202. Blue=ORF Red=Cloning site Green=Tag(s)

```

GCTCGTTTAGTGAACCGTCAGAATTTTGTAAACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTG
GATCCGGTACCGAGGAGATCTGCCGCCGCGATCGCC
ATGCCCGGCGAAGCCACAGAAACCGTCCCTGCTACAGAGCAGGAGTTGCCGCAGCCCCAGGCTGAGACA
GGGTCTGGAACAGAATCTGACAGTGATGAATCAGTACCAGAGCTTGAAGAACAGGATTCCACCCAGGCA
ACCACACAACAAGCCCAGCTGGCGGCAGCAGCTGAAATCGATGAAGAACCAGTCAGTAAAGCAAAACAG
AGTCGGAGTGAAAAGAAGGCACGGAAGGCTATGTCCAACTGGGTCTTCGGCAGGTTACAGGAGTTACT
AGAGTCACTATCCGAAATCTAAGAATATCCTCTTTGTCATCACAAAACCAGATGTCTACAAGAGCCCT
GCTTCAGATACTTACATAGTTTTTGGGGAAGCCAAGATCGAAGATTTATCCAGCAAGCACAACCTAGCA
GCTGCTGAGAAATTCAAAGTTCAAGGTGAAGCTGTCTCAAACATTCAAGAAAACACAGACTCCAAC
GTACAAGAGGAGAGTGAAGAGGAAGAGGTCGATGAAACAGGTGTAGAAGTTAAGGACATAGAATTGGTC
ATGTCACAAGCAAATGTGTGAGAGCAAAGGCAGTCCGAGCCCTGAAGAACAACAGTAATGATATTGTA
AATGCGATTATGGAATTAACAATG
ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGAT
TACAAGGATGACGACGATAAGGTTTAAACGGCCGCGC
  
```


[View online »](#)

Protein Sequence: >Peptide sequence encoded by RC225257
 Blue=ORF Red=Cloning site Green=Tag(s)

MPGEATETVPATEQELPQPPAETGSGTESDSDESVPLEEQDSTQATTQQAQLAAAAEIDEEPVSKAKQ
 SRSEKKARKAMSKLGLRQVTGVTRVTIRKSKNLFVITKPDVYKSPASDTYIVFGEAKIEDLSQQAQLA
 AAEKFKVQGEAVSNIQENTQPTVQEESEEEVDETGVEVKDIELVMSQANVSRKAVRALKNNSNDIV
 NAIMELTM
 TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Recombinant protein using RC225257 also available, [TP325257M](#)

Chromatograms: https://cdn.origene.com/chromatograms/mk6369_b01.zip

Restriction Sites: SgfI-MluI

Cloning Scheme:



ACCN: NM_001113202

ORF Size: 645 bp


OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.


Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:	1. Centrifuge at 5,000xg for 5min.
	2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
	3. Close the tube and incubate for 10 minutes at room temperature.
	4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
	5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
Note:	Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.
RefSeq ORF:	648 bp
Locus ID:	4666
UniProt ID:	Q13765
Cytogenetics:	12q13.3
Protein Families:	Druggable Genome, Transcription Factors
MW:	23.4 kDa
Gene Summary:	This gene encodes a protein that associates with basic transcription factor 3 (BTF3) to form the nascent polypeptide-associated complex (NAC). This complex binds to nascent proteins that lack a signal peptide motif as they emerge from the ribosome, blocking interaction with the signal recognition particle (SRP) and preventing mistranslocation to the endoplasmic reticulum. This protein is an IgE autoantigen in atopic dermatitis patients. Alternative splicing results in multiple transcript variants, but the full length nature of some of these variants, including those encoding very large proteins, has not been determined. There are multiple pseudogenes of this gene on different chromosomes. [provided by RefSeq, Feb 2016]


Product images:



Circular map for RC225257



Western blot validation of overexpression lysate (Cat# [LY426388]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with [RC225258] using transfection reagent MegaTran 2.0 (Cat# [TT210002]).



Coomassie blue staining of purified NACA protein (Cat# [TP325257]). The protein was produced from HEK293T cells transfected with NACA cDNA clone (Cat# RC225257) using MegaTran 2.0 (Cat# [TT210002]).