

Product datasheet for RC225345L2V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Bcl2 Binding component 3 (BBC3) (NM_001127240) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Bcl2 Binding component 3 (BBC3) (NM_001127240) Human Tagged ORF Clone Lentiviral

Particle

Symbol: Bcl2 Binding component 3

Synonyms: JFY-1; JFY1; PUMA

Mammalian Cell

Selection:

None

Vector: pLenti-C-mGFP (PS100071)

Tag: mGFP

ACCN: NM 001127240

ORF Size: 783 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC225345).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 001127240.1</u>

 RefSeq ORF:
 786 bp

 Locus ID:
 27113

 UniProt ID:
 Q9BXH1

 Cytogenetics:
 19q13.32

Protein Families: Druggable Genome

Protein Pathways: Huntington's disease, p53 signaling pathway



MW: 26.3 kDa

Gene Summary: This gene encodes a member of the BCL-2 family of proteins. This family member belongs to

the BH3-only pro-apoptotic subclass. The protein cooperates with direct activator proteins to induce mitochondrial outer membrane permeabilization and apoptosis. It can bind to anti-apoptotic Bcl-2 family members to induce mitochondrial dysfunction and caspase activation. Because of its pro-apoptotic role, this gene is a potential drug target for cancer therapy and for tissue injury. Alternative splicing results in multiple transcript variants. [provided by

RefSeq, Dec 2011]