

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for RC225711L1V

BLNK (NM_001114094) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	BLNK (NM_001114094) Human Tagged ORF Clone Lentiviral Particle
Symbol:	BLNK
Synonyms:	AGM4; BASH; bca; BLNK-S; LY57; SLP-65; SLP65
Mammalian Cell Selection:	None
Vector:	pLenti-C-Myc-DDK (PS100064)
Tag:	Myc-DDK
ACCN:	NM_001114094
ORF Size:	1299 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC225711).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 001114094.1, NP 001107566.1</u>
RefSeq ORF:	1302 bp
Locus ID:	29760
UniProt ID:	<u>Q8WV28</u>
Cytogenetics:	10q24.1
Protein Families:	Druggable Genome
Protein Pathways:	B cell receptor signaling pathway, Primary immunodeficiency
MW:	48 kDa



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



BLNK (NM_001114094) Human Tagged ORF Clone Lentiviral Particle – RC225711L1V

Gene Summary:This gene encodes a cytoplasmic linker or adaptor protein that plays a critical role in B cell
development. This protein bridges B cell receptor-associated kinase activation with
downstream signaling pathways, thereby affecting various biological functions. The
phosphorylation of five tyrosine residues is necessary for this protein to nucleate distinct
signaling effectors following B cell receptor activation. Mutations in this gene cause
hypoglobulinemia and absent B cells, a disease in which the pro- to pre-B-cell transition is
developmentally blocked. Deficiency in this protein has also been shown in some cases of
pre-B acute lymphoblastic leukemia. Alternatively spliced transcript variants have been found
for this gene. [provided by RefSeq, May 2012]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US