

## Product datasheet for RC225711L2

**OriGene Technologies, Inc.**  
 9620 Medical Center Drive, Ste 200  
 Rockville, MD 20850, US  
 Phone: +1-888-267-4436  
<https://www.origene.com>  
[techsupport@origene.com](mailto:techsupport@origene.com)  
 EU: [info-de@origene.com](mailto:info-de@origene.com)  
 CN: [techsupport@origene.cn](mailto:techsupport@origene.cn)

### BLNK (NM\_001114094) Human Tagged Lenti ORF Clone

#### Product data:



**Product Type:** Expression Plasmids  
**Tag:** mGFP  
**Symbol:** BLNK  
**Synonyms:** AGM4; BASH; bca; BLNK-S; LY57; SLP-65; SLP65  
**Mammalian Cell:** None  
**Selection:**  
**Vector:** pLenti-C-mGFP (PS100071)  
**E. coli Selection:** Chloramphenicol (34 ug/mL)

**ORF Nucleotide Sequence:** The ORF insert of this clone is exactly the same as (RC225711).

**Restriction Sites:** Sgfl-Mlul

#### Cloning Scheme:

Cloning sites used for ORF Shutting:



\* The last codon before the Stop codon of the ORF.

**ACCN:** NM\_001114094

**ORF Size:** 1299 bp



View online »

This product is to be used for laboratory only. Not for diagnostic or therapeutic use.

©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

<b>OTI Disclaimer:</b>	Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in <i>E. coli</i> are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at <a href="mailto:custsupport@origene.com">custsupport@origene.com</a> or by calling 301.340.3188 option 3 for pricing and delivery.
	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
<b>OTI Annotation:</b>	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>Note:</b>	Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.
<b>RefSeq:</b>	<a href="#">NM_001114094.1</a> , <a href="#">NP_001107566.1</a>
<b>RefSeq ORF:</b>	1302 bp
<b>Locus ID:</b>	29760
<b>UniProt ID:</b>	<a href="#">Q8WV28</a>
<b>Cytogenetics:</b>	10q24.1
<b>Protein Families:</b>	Druggable Genome
<b>Protein Pathways:</b>	B cell receptor signaling pathway, Primary immunodeficiency
<b>MW:</b>	48 kDa

**Gene Summary:**

This gene encodes a cytoplasmic linker or adaptor protein that plays a critical role in B cell development. This protein bridges B cell receptor-associated kinase activation with downstream signaling pathways, thereby affecting various biological functions. The phosphorylation of five tyrosine residues is necessary for this protein to nucleate distinct signaling effectors following B cell receptor activation. Mutations in this gene cause hypoglobulinemia and absent B cells, a disease in which the pro- to pre-B-cell transition is developmentally blocked. Deficiency in this protein has also been shown in some cases of pre-B acute lymphoblastic leukemia. Alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, May 2012]

**Product images:**