

Product datasheet for RC225740L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CSK (NM_001127190) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: CSK (NM_001127190) Human Tagged ORF Clone Lentiviral Particle

Symbol: CSK

Mammalian Cell Puromycin

Selection:

Vector:

pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK

ACCN: NM_001127190

ORF Size: 1350 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC225740).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 001127190.1</u>

 RefSeq Size:
 2236 bp

 RefSeq ORF:
 1353 bp

 Locus ID:
 1445

 UniProt ID:
 P41240

 Cytogenetics:
 15q24.1

Protein Families: Druggable Genome, Protein Kinase

Protein Pathways: Chemokine signaling pathway, Epithelial cell signaling in Helicobacter pylori infection,

Neurotrophin signaling pathway, Regulation of actin cytoskeleton







MW:

50.7 kDa

Gene Summary:

The protein encoded by this gene is involved in multiple pathways, including the regulation of Src family kinases. It plays an important role in T-cell activation through its association with the protein encoded by the protein tyrosine phosphatase, non-receptor type 22 (PTPN22) gene. This protein also phosphorylates C-terminal tyrosine residues on multiple substrates, including the protein encoded by the SRC proto-oncogene, non-receptor tyrosine kinase gene. Phosphorylation suppresses the kinase activity of the Src family tyrosine kinases. An intronic polymorphism (rs34933034) in this gene has been found to affect B-cell activation and is associated with systemic lupus erythematosus (SLE). Alternative splicing results in multiple transcript variants. [provided by RefSeq, Aug 2017]