

### Product datasheet for RC225775L2

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## PI 3 Kinase p55 gamma (PIK3R3) (NM\_001114172) Human Tagged Lenti ORF Clone

#### **Product data:**

**Product Type:** Expression Plasmids

Product Name: PI 3 Kinase p55 gamma (PIK3R3) (NM\_001114172) Human Tagged Lenti ORF Clone

Tag: mGFP

Symbol: PI 3 Kinase p55 gamma

**Synonyms:** p55; p55-GAMMA; p55PIK

Mammalian Cell None

Selection:

**Vector:** pLenti-C-mGFP (PS100071)

E. coli Selection: Chloramphenicol (34 ug/mL)


**ORF Nucleotide** The ORF insert of this clone is exactly the same as(RC225775).

Sequence:

Restriction Sites: Sgfl-Mlul

Cloning Scheme:





<sup>\*</sup> The last codon before the Stop codon of the ORF.

**ACCN:** NM\_001114172

ORF Size: 1383 bp



ORIGENE

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001114172.1</u>, <u>NP 001107644.1</u>

 RefSeq ORF:
 1386 bp

 Locus ID:
 8503

 UniProt ID:
 Q92569

 Cytogenetics:
 1p34.1

**Protein Families:** Druggable Genome

**Protein Pathways:** Acute myeloid leukemia, Apoptosis, B cell receptor signaling pathway, Chemokine signaling

pathway, Chronic myeloid leukemia, Colorectal cancer, Endometrial cancer, ErbB signaling pathway, Fc epsilon RI signaling pathway, Fc gamma R-mediated phagocytosis, Focal adhesion, Glioma, Insulin signaling pathway, Jak-STAT signaling pathway, Leukocyte

transendothelial migration, Melanoma, mTOR signaling pathway, Natural killer cell mediated cytotoxicity, Neurotrophin signaling pathway, Non-small cell lung cancer, Pancreatic cancer, Pathways in cancer, Phosphatidylinositol signaling system, Progesterone-mediated oocyte maturation, Prostate cancer, Regulation of actin cytoskeleton, Renal cell carcinoma, Small cell lung cancer, T cell receptor signaling pathway, Toll-like receptor signaling pathway, Type II

diabetes mellitus, VEGF signaling pathway

**MW:** 54.3 kDa

**Gene Summary:** Phosphatidylinositol 3-kinase (PI3K) phosphorylates phosphatidylinositol and similar


compounds, which then serve as second messengers in growth signaling pathways. PI3K is composed of a catalytic and a regulatory subunit. The protein encoded by this gene represents a regulatory subunit of PI3K. The encoded protein contains two SH2 domains

through which it binds activated protein tyrosine kinases to regulate their activity. [provided

by RefSeq, Jun 2016]



# **Product images:**



Circular map for RC225775L2