

# **Product datasheet for RC225964L2**

## **OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

## WEE2 (NM\_001105558) Human Tagged Lenti ORF Clone

#### **Product data:**

**Product Type:** Expression Plasmids

**Product Name:** WEE2 (NM\_001105558) Human Tagged Lenti ORF Clone

Tag: mGFP Symbol: WEE2

Synonyms: OOMD5; WEE1B

Mammalian Cell None

Selection:

**Vector:** pLenti-C-mGFP (PS100071)

E. coli Selection: Chloramphenicol (34 ug/mL)

ORF Nucleotide The ORF insert of this clone is exactly the same as(RC225964).

Sequence:

**Restriction Sites:** Sgfl-Mlul

**Cloning Scheme:** 





<sup>\*</sup> The last codon before the Stop codon of the ORF.

**ACCN:** NM\_001105558

ORF Size: 1701 bp





### WEE2 (NM\_001105558) Human Tagged Lenti ORF Clone - RC225964L2

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

**RefSeq:** <u>NM 001105558.1, NP 001099028.1</u>

**RefSeq ORF:** 1704 bp **Locus ID:** 494551

UniProt ID: P0C1S8

Cytogenetics: 7q34

Protein Pathways: Cell cycle

**MW:** 62.7 kDa


Gene Summary: Oocyte-specific protein tyrosine kinase that phosphorylates and inhibits CDK1/CDC2 and acts

as a key regulator of meiosis during both prophase I and metaphase II (PubMed:29606300). Required to maintain meiotic arrest in oocytes during the germinal vesicle (GV) stage, a long period of quiescence at dictyate prophase I, by phosphorylating CDK1 at 'Tyr-15', leading to inhibit CDK1 activity and prevent meiotic reentry. Also required for metaphase II exit during egg activation by phosphorylating CDK1 at 'Tyr-15', to ensure exit from meiosis in oocytes and

promote pronuclear formation (By similarity).[UniProtKB/Swiss-Prot Function]



# **Product images:**



Circular map for RC225964L2