

## Product datasheet for RC225984L1V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## GAD65 (GAD2) (NM 001134366) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** GAD65 (GAD2) (NM\_001134366) Human Tagged ORF Clone Lentiviral Particle

Symbol: GAD65 Synonyms: **Mammalian Cell** 

Selection:

None

Vector: pLenti-C-Myc-DDK (PS100064)

Myc-DDK Tag:

NM 001134366 ACCN:

**ORF Size:** 1755 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC225984).

Sequence: OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 001134366.1

RefSeq Size: 2419 bp RefSeq ORF: 1758 bp Locus ID: 2572 **UniProt ID:** Q05329 Cytogenetics: 10p12.1

**Protein Families:** Druggable Genome





## GAD65 (GAD2) (NM\_001134366) Human Tagged ORF Clone Lentiviral Particle - RC225984L1V

**Protein Pathways:** Alanine, aspartate and glutamate metabolism, beta-Alanine metabolism, Butanoate

metabolism, Metabolic pathways, Taurine and hypotaurine metabolism, Type I diabetes

mellitus

MW: 65.4 kDa

**Gene Summary:** This gene encodes one of several forms of glutamic acid decarboxylase, identified as a major

autoantigen in insulin-dependent diabetes. The enzyme encoded is responsible for catalyzing the production of gamma-aminobutyric acid from L-glutamic acid. A pathogenic role for this enzyme has been identified in the human pancreas since it has been identified as an autoantibody and an autoreactive T cell target in insulin-dependent diabetes. This gene may also play a role in the stiff man syndrome. Alternative splicing results in multiple transcript

variants that encode the same protein. [provided by RefSeq, Oct 2008]