

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for RC226000L1V

NKIAMRE (CDKL3) (NM_001113575) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	NKIAMRE (CDKL3) (NM_001113575) Human Tagged ORF Clone Lentiviral Particle
Symbol:	NKIAMRE
Synonyms:	NKIAMRE
Mammalian Cell Selection:	None
Vector:	pLenti-C-Myc-DDK (PS100064)
Tag:	Myc-DDK
ACCN:	NM_001113575
ORF Size:	1776 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC226000).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 001113575.1</u>
RefSeq ORF:	1779 bp
Locus ID:	51265
UniProt ID:	<u>Q8IVW4</u>
Cytogenetics:	5q31.1
Protein Families:	Druggable Genome, Protein Kinase
MW:	67.3 kDa



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Gene Summary:The protein encoded by this gene is a member of cyclin-dependent protein kinase (CDK)
family. CDK family members are highly similar to the gene products of Saccharomyces
cerevisiae cdc28, and Schizosaccharomyces pombe cdc2, and are known to be important
regulators of cell cycle progression. This gene was identified as a gene absent in leukemic
patients with chromosome 5q deletion. This loss may be an important determinant of
dysmyelopoiesis. Alternative splicing results in multiple transcript variants encoding different
isoforms. [provided by RefSeq, Jul 2008]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US