

## Product datasheet for RC226044L1V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## SHTN1 (NM\_001127211) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** SHTN1 (NM\_001127211) Human Tagged ORF Clone Lentiviral Particle

Symbol: SHTN1

Synonyms: KIAA1598; shootin-1

Mammalian Cell

Selection:

None

**Vector:** pLenti-C-Myc-DDK (PS100064)

Tag: Myc-DDK

**ACCN:** NM\_001127211

ORF Size: 1893 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC226044).

Sequence:

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 001127211.1</u>

 RefSeq ORF:
 1896 bp

 Locus ID:
 57698

 UniProt ID:
 A0MZ66

 Cytogenetics:
 10q25.3

**MW:** 71.5 kDa





## **Gene Summary:**

Involved in the generation of internal asymmetric signals required for neuronal polarization and neurite outgrowth. Mediates netrin-1-induced F-actin-substrate coupling or 'clutch engagement' within the axon growth cone through activation of CDC42, RAC1 and PAK1-dependent signaling pathway, thereby converting the F-actin retrograde flow into traction forces, concomitantly with filopodium extension and axon outgrowth. Plays a role in cytoskeletal organization by regulating the subcellular localization of phosphoinositide 3-kinase (PI3K) activity at the axonal growth cone. Plays also a role in regenerative neurite outgrowth. In the developing cortex, cooperates with KIF20B to promote both the transition from the multipolar to the bipolar stage and the radial migration of cortical neurons from the ventricular zone toward the superficial layer of the neocortex. Involved in the accumulation of phosphatidylinositol 3,4,5-trisphosphate (PIP3) in the growth cone of primary hippocampal neurons.[UniProtKB/Swiss-Prot Function]