

Product datasheet for **RC226067**

PEX5 (NM_001131023) Human Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	PEX5 (NM_001131023) Human Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	PEX5
Synonyms:	PBD2A; PBD2B; PTS1-BP; PTS1R; PXR1; RCDP5
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)



[View online »](#)

ORF Nucleotide
Sequence:

>RC226067 representing NM_001131023
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGATCGCC**

ATGGCAATGCGGGAGCTGGTGGAGGCCAATGCGGGGTGCCAACCCGCTCATGAAGCTCGCCGGGCACT
 TCACCCAGGACAAGGCCCTTCGGCAGGAGGGATTGAGGCCTGGCCCTGGCCCCGGAGCCCCGGCCTC
 TGAGGCAGTGAGTGTTCTTGAGGTGGAAGCCAGGTGCAGCCTCTGAGGCAGCCTCCAAGCCTTTGGGA
 GTAGTCTCTGAAGATGAGTTGGTGGCTGAATTCCTGCAGGACCAGAATGCACCCCTTGTTGCCGTGCC
 CTCAGACCTTCAAGATGGATGACCTCCTGGCTGAGATGCAGCAGATTGAGCAGTCAAACCTCCGCCAGGC
 TCCCCAGAGAGCCCTGGTGTGGCAGACTTGGCCTTGTCTGAGAAGTGGGCCAGGAGTTTCTTGACGCT
 GGAGATGCTGTGGATGTAACCTCAGGATTATAATGAGACTGACTGGTCCCAAGAATTCATCTCTGAAGTTA
 CAGACCCCTTGTCTGTGTCCCCTGCCCGTGGGCTGAGGAATATTTGGAGCAATCAGAGGAGAAGCTGTG
 GCTGGGAGAACCAGGGAAACAGCCACCGATCGCTGGTATGATGAATATCATCTGAGGAGGATCTGCAG
 CACACGGCCAGTGACTTTGTGGCCAAAGTGGATGACCCCAAATGGCTAATTCAGTTCCTGAAATTCG
 TGCGGCAGATTGGCGAAGGGCAGGTGTCCCTGGAGTCCGGTGCAGGGTCCGGCCGAGCTCAGGCAGAACA
 GTGGGCAGCAGAGTTTATACAGCAGCAGGGTACATCAGATGCCTGGGTTGACCAGTTCACAAGACCAGTA
 AACACATCTGCCCTTGATATGGAGTTTGAACGAGCCAAGTCAGCTATAGAGTCTGATGTCGATTTCTGGG
 ACAAGTTGCAGGCAGAGTTGGAGGAGATGGCAAAACGGGATGCTGAGGCCACCCCTGGCTTTCTGACTA
 TGATGACCTTACGTCAGTACCTATGATAAGGGGTACCAGTTTGGAGGAGAGAACCCTTGGCGTATCAC
 CCTCAGCCTTTTGAAGAAGGGCTGCGGCCCTTCAGGAGGGGGACCTGCCAAATGCTGTGCTGTTTTTG
 AGGCAGCTGTGCAGCAGGATCCTAAGCACATGGAAGCTTGGCAGTATCTGGGTACCACCCAGGCAGAGAA
 TGAACAAGAACTATTAGCCATCAGTGCATTGCGGAGGTGTCTGGAGCTAAAGCCAGATAACCAGACAGCA
 CTGATGGCGCTGGCTGTGAGCTTACCAACGAGTCCCTGCAGCGACAGGCCTGTGAAACCTACGAGACT
 GGCTGCGGTACACACCAGCCTATGCCCATCTGGTGACACCTGCTGAAGAAGGGGCTGGTGGGGCAGGACT
 GGGCCCCAGCAAGCGTATCCTGGGATCTCTTGTCTGACTCCCTGTTTCTTGAAGTGAAGAGCTCTTC
 CTGGCAGCTGTGCGGCTGGACCTACCTCCATTGACCCTGATGTGCAGTGTGGCTTGGGAGTCTTTTCA
 ACCTGAGTGGGGAGTATGACAAGGCCGTGGACTGCTTACAGCTGCCCTCAGCGTTCGTCCCAATGACTA
 TTTGCTGTGGAATAAGCTAGGCGCCACCCTGGCCAATGGAACCAGAGTGAAGAAGCAGTAGCTGCGTAC
 CGCCGGGCCCTCGAGCTCCAGCCTGGCTATATCCGGTCCCCTATAACCTGGGCATCAGCTGCATCAACC
 TCGGGGCTCACCGGAGGCTGTGGAGCACTTTCTGGAGGCCCTGAACATGCAGAGGAAAAGCCGGGGCCC
 CCGGGGTGAAGGAGGTGCCATGTGCGGAGAACATCTGGAGCACCTGCGTTTGGCATTGTCTATGTTAGGC
 CAGAGCGATGCCTATGGGGCAGCCGACGCGGGGATCTGTCCACCCTCTAACTATGTTTGGCCTGCCCC
 AG

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence: >RC226067 representing NM_001131023
 Red=Cloning site Green=Tags(s)

MAMRELVEAECGGANPLMKLAGHFTQDKALRQEGLRPGWPPGAPASEAVSVLEVESPGAASEAASKPLG
 VASEDELVAEFLQDNAPLVSRAPQTFKMDDLAEQQIEQSNFRQAPQAPGVADLALSENWAQEFLLA
 GDAVDVTQDYNETDWSQEFISEVTDPLSVSPARWAEYLEQSEEKWLGEPEGTATDRWYDEYHPEEDLQ
 HTASDFVAKVDDPKLANSEFLKFVRQIGEGQVLSLESGAGSGRAQAEQWAAEFIQQQGTSDAWVDQFTRPV
 NTSALDMEFERAKSAIESDVFWDKLAELLEEMAKRDAEAHPWLSYDDDLTSATYDKGYQFEEENPLRDH
 PQPFEEGLRRLQEGDLPNAVLLFEAAVQQDPKHMEAWQYLGTQAENEQELLAISALRRCLELKPNDQTA
 LMALAVSFTNESLQRQACETLRDWLRYTPAYAHLVTPAEEGAGGAGLGPSKRILGSLSDSLFLEVKELF
 LAAVRLDPTSIDPDVQCGLGVLFNLSGEYDKAVDCFTAALSVRPNDYLLWNKLGATLANGQSEEAVAA
 RRALQLPGYIRSRYNLGISCNLGAHREAVEHFLEALNMQRKSRGPRGEGGAMSENIWSTLRLALSMGL
 QSDAYGAADARDLSTLLTMFGLPQ


TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Chromatograms: https://cdn.origene.com/chromatograms/mk8071_f05.zip

Restriction Sites: SgfI-MluI

Cloning Scheme:

Cloning sites used for ORF Shuttling:



* The last codon before the Stop codon of the ORF

ACCN: NM_001131023

ORF Size: 1962 bp

OTI Disclaimer: Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at custsupport@origene.com or by calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001131023.2](#)

RefSeq ORF: 1965 bp

Locus ID: 5830

UniProt ID: [P50542](#)

Cytogenetics: 12p13.31


Protein Families: Druggable Genome

MW: 72.1 kDa


Gene Summary:

The product of this gene binds to the C-terminal PTS1-type tripeptide peroxisomal targeting signal (SKL-type) and plays an essential role in peroxisomal protein import. Peroxins (PEXs) are proteins that are essential for the assembly of functional peroxisomes. The peroxisome biogenesis disorders (PBDs) are a group of genetically heterogeneous autosomal recessive, lethal diseases characterized by multiple defects in peroxisome function. The peroxisomal biogenesis disorders are a heterogeneous group with at least 14 complementation groups and with more than 1 phenotype being observed in cases falling into particular complementation groups. Although the clinical features of PBD patients vary, cells from all PBD patients exhibit a defect in the import of one or more classes of peroxisomal matrix proteins into the organelle. Defects in this gene are a cause of neonatal adrenoleukodystrophy (NALD), a cause of Zellweger syndrome (ZWS) as well as may be a cause of infantile Refsum disease (IRD). Alternatively spliced transcript variants encoding different isoforms have been identified. [provided by RefSeq, Oct 2008]

Product images:



Circular map for RC226067



Western blot validation of overexpression lysate (Cat# [LY427361]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with RC226067 using transfection reagent MegaTran 2.0 (Cat# [TT210002]).