

## Product datasheet for **RC226264**

### **DIS3 (NM\_001128226) Human Tagged ORF Clone**

#### **Product data:**

|                           |  |
|---------------------------|--|
| Product Type:             | Expression Plasmids                            |
| Product Name:             | DIS3 (NM_001128226) Human Tagged ORF Clone     |
| Tag:                      | Myc-DDK  |
| Symbol:                   | DIS3   |
| Synonyms:                 | 2810028N01Rik; dis3p; EXOSC11; KIAA1008; RRP44 |
| Mammalian Cell Selection: | Neomycin                                       |
| Vector:                   | pCMV6-Entry (PS100001)                         |
| E. coli Selection:        | Kanamycin (25 ug/mL)                           |



[View online »](#)

ORF Nucleotide  
Sequence:

>RC226264 representing NM\_001128226  
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
 GCC**CGGATCGCC**


ATGCTCAAGTCCAAGACGTTCTTAAAAAGACCCGGGCGGGCGGCGTGATGAAGATCGTGCAGCACT  
 ACCTGCGAGACGACATCGCTGCGGTGCGCCCGGTGCGCAGCGTGTGGAGGGCGCACGAGGGCCGGC  
 CCTGGAGCCGAGCCCCAGGACCCGCGAGCAGCGTCTGCCCGCAACCGCACTACTTGTGCCGCACT  
 AATGTGTTACTGCACCAGATTGTAAGTGCTGGAGGCCGGGACCTGGGTTCTGTGGCCTCCAGCCTGC  
 GACTCCCAGGAGCTTAGAACTATGTAGAACAAGAAGAGGAGAAAATGCTAATGACAGGAATGATAG  
 AGCGATTCGAGTAGCAGCAAAATGGTACAATGAACATTTGAAAAAATGTCAGCAGACAACCAGCTGCAA  
 GTTATCTTCATAACAAATGACAGGAGAAAACAAGAGAAAAGCCATAGAAGAAGGAATACCAGCTTTCCTT  
 GTGAAGAATATGTAAGAGCCTAACTGCTAACCCCGAAGCTCATAGATCGTCTTGTGTTGTCTGAAGA  
 AGGGAATGAAATAGAAAGTGGAAAAATAATATTTTCAGAGCATCTCCCTTAAGTAAGCTACAGCAAGGC  
 ATAAAATCTGGTACATACCTTCAAGGAACATTTAGAGCTAGCAGGGAAAATTAAGTGAAGCTACAGCAAGGC  
 GGATTCATGGCGACAATGAAGAAAATAAAGAGATAATCTTACAGGGACTTAAACATTTAAACAGAGCTGT  
 TCACGAAGATATTGTGGCTGTGGAGCTTCTCCCAAGAGTCAGTGGGTAGCACCATCTTCTGTGGTTTTA  
 CATGATGAAGGTCAAAATGAAGAAGATGTGGAGAAAGAAGAAGAGACAGAACGAATGCTTAAAGACTGCTG  
 TAAGCGAGAAAATGTTGAAGCCTACAGGTAGAGTTGTAGGAATAATAAAAGGAATTGGAGACCATTG  
 TGGCATGCTTCCAAGTCTGACATTAAGGAGTCAAGAAGACATCTTTACACCTGCTGATAAGAGAATC  
 CCTCGAATTCGCATAGAAACCAGACAGGCTTCCACATTAGAAGGACGGAGAATTATTGTTGCTATTGATG  
 GTTGGCCAGAAAATCCAGATATCCAAATGGACACTTTGTGAGAAAATTTAGGTGATGTTGGAGAGAAAAG  
 GACTGAAACAGAAAGTTTTGTTACTTGAACACGATGTTCCCATCAGCCTTTTTACAGGCTGTTCTTAGT  
 TTTCTGCCAAAGATGCCCTGGAGCATTACTGAAAAGGACATGAAAAACCGAGAAGACCTGAGGCATCTGT  
 GTATTTGTAGTGTAGACCACCAGGATGTACTGATATAGACGATGCTCTACATTGTGCGAACTCGAAAA  
 TGGAAATTTGGAGGTTGGTTCATATTGCTGATGTGAGCCATTTTATTAGGCCAGGAAATGCCTTGGAT  
 CAAGAATCAGCCAGAAGAGGAACAACCTGTGATCTTTGTGAAAAGAGGATTGACATGGTTCCAGAGTTGC  
 TTAGCTCTAACTTGTGTTCTTAAAATGTGACGTGGACAGGCTGGCATTTCATGATTTGGGAAATGAA  
 TCACAATGCTGAAATCTTAAAAACGAAGTTTACCAAAAGTGTATTAATCAAAGGCATCTCTGACGTAT  
 GCTGAAGCTCAGTTGAGAATTGATTCAGCAAACATGAATGATGATATTACCACTAGTCTCCGTGGACTGA  
 ATAAACTAGCCAAAATCTGAAGAAAAGAAGGATTGAAAAAGGGCTTTGACTCTATCCTCTCCTGAAGT  
 TCGATTCCACATGGACAGTGAAGCTCACGATCCTATAGATCTGCAGACCAAGGAAGTGGGAAACAAAT  
 TCCATGGTTGAAGAAATTTATGTTACTTGCCAATATTTCTGTTGCAAAAAAATTCATGAGGAATTTCTG  
 AACATGCTCTGCTCGAAAACATCCTGCTCCACCTCCATCAAAATTAAGAAATTTGTTAAGGCAGCCAG  
 GTCAAGGAATTTGGAAATTAAGACTGATACAGCCAAGTCTTTGGCTGAGTCTTTGGATCAGGCCGAATCT  
 CCTACTTTCCATATCTAAACACTCTGTTGAGAATATTAGCCACTCGCTGATGATGCAAGCTGTGACT  
 TCTGTTCTGGAATGGATAATGATTTTCATCACTATGGCTTAGCGTCTCCAATATACACACATTTACTTC  
 ACCCATTAGAAGATACGCAGATGTCATTGTTTCATCGGCTTTTGGCTGTGGCTATTGGGGCTGACTGTACT  
 TATCCAGAGTTGACAGACAACACAAGCTTGAGATATATGAAAAATCTAAATTTCCGGCACAAAATGG  
 CTCAATATGCCAACGTGCATCAGTGGCTTTTCATACCCAGTTATTCTTCAAAGCAAAGGAATAGTAAG  
 TGAAGAAGCCTATATTTATTTGTAAGAAAAGATGCCATTGTGGTATTAATCCAAAGTATGGTTTAGAA  
 GGGACAGTCTTTTTGAAAGAAAAGGACAAACCAACCCACAGCTTATTTATGATGATGAGATACCCTCAC  
 TAAAAATAGAAGATACAGTGTCCATGTATTTGATAAAGTTAAAGTAAAATCATGTTAGACTCATCTAA  
 TCTTCAACATCAGAAGATCCGAATGTCCCTGGTAGAACACAGATACCAGGAATAAGCATTCTACTGAT  
 ACTTCAAACATGGACCTAATGGACCAAGAAAAGAAGATGAAGCTTGGAAAA

**ACGCGT**ACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT  
 ACAAGGATGACGACGATAAAGTTTAA



|                               |  |
|-------------------------------|--|
| <b>OTI Disclaimer:</b>        | The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>   |
| <b>OTI Annotation:</b>        | This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.   |
| <b>Components:</b>            | The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).   |
| <b>Reconstitution Method:</b> | <ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>  |
| <b>RefSeq:</b>                | <a href="#">NM_001128226.3</a>   |
| <b>RefSeq ORF:</b>            | 2787 bp  |
| <b>Locus ID:</b>              | 22894  |
| <b>UniProt ID:</b>            | <a href="#">Q9Y2L1</a>   |
| <b>Cytogenetics:</b>          | 13q21.33   |
| <b>Protein Pathways:</b>      | RNA degradation  |
| <b>MW:</b>                    | 104.9 kDa  |
| <b>Gene Summary:</b>          | Putative catalytic component of the RNA exosome complex which has 3'->5' exoribonuclease activity and participates in a multitude of cellular RNA processing and degradation events. In the nucleus, the RNA exosome complex is involved in proper maturation of stable RNA species such as rRNA, snRNA and snoRNA, in the elimination of RNA processing by-products and non-coding 'pervasive' transcripts, such as antisense RNA species and promoter-upstream transcripts (PROMPTs), and of mRNAs with processing defects, thereby limiting or excluding their export to the cytoplasm. The RNA exosome may be involved in Ig class switch recombination (CSR) and/or Ig variable region somatic hypermutation (SHM) by targeting AICDA deamination activity to transcribed dsDNA substrates. In the cytoplasm, the RNA exosome complex is involved in general mRNA turnover and specifically degrades inherently unstable mRNAs containing AU-rich elements (AREs) within their 3' untranslated regions, and in RNA surveillance pathways, preventing translation of aberrant mRNAs. It seems to be involved in degradation of histone mRNA. DIS3 has both 3'-5' exonuclease and endonuclease activities.[UniProtKB/Swiss-Prot Function] |

Product images:



Circular map for RC226264