

## **Product datasheet for RC226296L3**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### MOV10 (NM\_001130079) Human Tagged Lenti ORF Clone

#### **Product data:**

**Product Type:** Expression Plasmids

**Product Name:** MOV10 (NM\_001130079) Human Tagged Lenti ORF Clone

Tag: Myc-DDK Symbol: MOV10

**Synonyms:** fSAP113; gb110

Mammalian Cell Puromycin

Selection:

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

E. coli Selection: Chloramphenicol (34 ug/mL)

ORF Nucleotide The ORF insert of this clone is exactly the same as(RC226296).

Sequence:

Restriction Sites: Sgfl-Mlul

**Cloning Scheme:** 





st The last codon before the Stop codon of the ORF.

**ACCN:** NM\_001130079

ORF Size: 3009 bp



#### MOV10 (NM\_001130079) Human Tagged Lenti ORF Clone - RC226296L3

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001130079.1</u>, <u>NP 001123551.1</u>

RefSeq Size: 3491 bp
RefSeq ORF: 3012 bp
Locus ID: 4343
UniProt ID: Q9HCE1

Cytogenetics: 1p13.2

**MW:** 113.7 kDa

**Gene Summary:** 5' to 3' RNA helicase contributing to UPF1 mRNA target degradation by translocation along 3'

UTRs (PubMed:24726324). Required for microRNA (miRNA)-mediated gene silencing by the RNA-induced silencing complex (RISC). Required for both miRNA-mediated translational

repression and miRNA-mediated cleavage of complementary mRNAs by RISC

(PubMed:16289642, PubMed:17507929, PubMed:22791714). In cooperation with FMR1, regulates miRNA-mediated translational repression by AGO2 (PubMed:25464849). Restricts retrotransposition of long interspersed element-1 (LINE-1) in cooperation with TUT4 and

TUT7 counteracting the RNA chaperonne activity of L1RE1 (PubMed:30122351,

PubMed:23093941). Facilitates LINE-1 uridylation by TUT4 and TUT7 (PubMed:30122351). Required for embryonic viability and for normal central nervous system development and function. Plays two critical roles in early brain development: suppresses retroelements in the nucleus by directly inhibiting cDNA synthesis, while regulates cytoskeletal mRNAs to influence


neurite outgrowth in the cytosol (By similarity). May function as a messenger

ribonucleoprotein (mRNP) clearance factor (PubMed:24726324).[UniProtKB/Swiss-Prot

Function]



# **Product images:**



Circular map for RC226296L3