

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for RC226319L3V

Angiomotin (AMOT) (NM_001113490) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	Angiomotin (AMOT) (NM_001113490) Human Tagged ORF Clone Lentiviral Particle
Symbol:	АМОТ
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_001113490
ORF Size:	3252 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC226319).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 001113490.1, NP 001106962.1</u>
RefSeq ORF:	3255 bp
Locus ID:	154796
UniProt ID:	Q4VCS5
Cytogenetics:	Xq23
MW:	117.9 kDa



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	Angiomotin (AMOT) (NM_001113490) Human Tagged ORF Clone Lentiviral Particle – RC226319L3V
Gene Summary:	This gene belongs to the motin family of angiostatin binding proteins characterized by
	conserved coiled-coil domains and C-terminal PDZ binding motifs. The encoded protein is

expressed predominantly in endothelial cells of capillaries as well as larger vessels of the placenta where it may mediate the inhibitory effect of angiostatin on tube formation and the migration of endothelial cells toward growth factors during the formation of new blood vessels. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Jul 2008]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US