

## Product datasheet for **RC226555L3V**

### 26S proteasome non ATPase regulatory subunit 12 (PSMD12) (NM\_174871) Human Tagged ORF Clone Lentiviral Particle

#### Product data:

|                           |  |
|---------------------------|--|
| Product Type:             | Lentiviral Particles   |
| Product Name:             | 26S proteasome non ATPase regulatory subunit 12 (PSMD12) (NM_174871) Human Tagged ORF Clone Lentiviral Particle  |
| Symbol:                   | 26S proteasome non ATPase regulatory subunit 12  |
| Synonyms:                 | p55; Rpn5; STISS   |
| Mammalian Cell Selection: | Puromycin  |
| Vector:                   | pLenti-C-Myc-DDK-P2A-Puro (PS100092)   |
| Tag:                      | Myc-DDK  |
| ACCN:                     | NM_174871  |
| ORF Size:                 | 1308 bp  |
| ORF Nucleotide Sequence:  | The ORF insert of this clone is exactly the same as(RC226555).   |
| OTI Disclaimer:           | <p>Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at <a href="mailto:custsupport@origene.com">custsupport@origene.com</a> or by calling 301.340.3188 option 3 for pricing and delivery.</p> <p>The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a></p> |
| OTI Annotation:           | This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.   |
| RefSeq:                   | <a href="#">NM_174871.2</a>  |



[View online »](#)

RefSeq ORF: 1311 bp

Locus ID: 5718

UniProt ID: [O00232](#)

Cytogenetics: 17q24.2

Protein Pathways: Proteasome

MW: 50.4 kDa

**Gene Summary:** The 26S proteasome is a multicatalytic proteinase complex with a highly ordered structure composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. The 19S regulator is composed of a base, which contains 6 ATPase subunits and 2 non-ATPase subunits, and a lid, which contains up to 10 non-ATPase subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a non-ATPase subunit of the 19S regulator. A pseudogene has been identified on chromosome 3. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Oct 2015]