

Product datasheet for RC226571L2V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

IL11RA (NM_001142784) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: IL11RA (NM_001142784) Human Tagged ORF Clone Lentiviral Particle

Symbol: IL11RA
Synonyms: CRSDA

Mammalian Cell

Selection:

None

Vector: pLenti-C-mGFP (PS100071)

Tag: mGFP

ACCN: NM_001142784

ORF Size: 1266 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC226571).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 001142784.1

RefSeq Size: 1728 bp
RefSeq ORF: 1269 bp
Locus ID: 3590
UniProt ID: Q14626
Cytogenetics: 9p13.3

Protein Families: Druggable Genome, Transmembrane





IL11RA (NM_001142784) Human Tagged ORF Clone Lentiviral Particle - RC226571L2V

Protein Pathways: Cytokine-cytokine receptor interaction, Hematopoietic cell lineage, Jak-STAT signaling

pathway

MW: 45.2 kDa

Gene Summary: Interleukin 11 is a stromal cell-derived cytokine that belongs to a family of pleiotropic and

redundant cytokines that use the gp130 transducing subunit in their high affinity receptors. This gene encodes the IL-11 receptor, which is a member of the hematopoietic cytokine receptor family. This particular receptor is very similar to ciliary neurotrophic factor, since

both contain an extracellular region with a 2-domain structure composed of an

immunoglobulin-like domain and a cytokine receptor-like domain. Multiple alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Jun 2012]