

Product datasheet for RC226612

NEIL2 (NM_001135748) Human Tagged ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: NEIL2 (NM_001135748) Human Tagged ORF Clone

Tag: Myc-DDK

Symbol: NEIL2

Synonyms: NEH2; NEI2

Mammalian Cell

Selection:

Neomycin

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

ORF Nucleotide >RC226612 representing NM_001135748
Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ATGCCAGAAGGGCCGTTGGTGAGGAAATTTCACCATTTGGTCTCCCCCTTTGTGGGTCAGCAGGTGGTCA
AGACAGGGGGCAGCAGCAGAAGAAGCTACAGCCCGCCAGCCTGCAGTCTCTGTGGCTCCAGGACACCCAGGT
GAGGTTGGTCCTGCACTTTGGTGGTGGTGGTCCTCTGGCATTTTATAATTGTCAGTTGTCTTGGAGCTCT
TCCCCAGTGGTCACACCCACCTGTGACATCCTGTCTGAGAAGTTCCATCGAGGACAAGCCTTAGAAGCTC
TAGGCCAGGCTCAGCCTGTCTGCTATACACTGCTGGACCAGAGATACTTCTCAGGGCTAGGGAACATCAT
TAAGAATGAAGCCTTGTACAGAGCTGGGATCCATCCCCTTTCTCTCTGGTTCAGTCCTGAGTGCCTCGCGT
CGGGAGGTCCTGGTGGATCACCTGGTGGAGTTCAGTACAGCCTGGCTGCAGGGCAAGTTCCAAGGCAGAC
CGCAGCACACACAGGTCTACCAGAAAGAACAGTGCCCTGCTGGCCACCAGGTCATGAAGGAGGCGTTTGG
GCCCGAAGATGGGTTACAGAGGCTCACCTGGTGGTGCCCCCAGTTGCCAGCCCCAGTTTCAGAGGAGCCA
GAGCAGTGCCAGTTCTCC

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATTACAAGGATGACGACGATAAGGTTTAA



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

NEIL2 (NM_001135748) Human Tagged ORF Clone - RC226612

Protein Sequence: >RC226612 representing NM_001135748

Red=Cloning site Green=Tags(s)

MPEGPLVRKFHHLVSPFVGQQVVKTGGSSKKLQPASLQSLWLQDTQVRLVLHFGGGGFLAFYNCQLSWSS SPVVTPTCDILSEKFHRGQALEALGQAQPVCYTLLDQRYFSGLGNIIKNEALYRAGIHPLSLGSVLSASR REVLVDHVVEFSTAWLQGKFQGRPQHTQVYQKEQCPAGHQVMKEAFGPEDGLQRLTWWCPQCQPQLSEEP EQCQFS


TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Chromatograms: https://cdn.origene.com/chromatograms/ja1433 c07.zip

Restriction Sites: Sgfl-Mlul

Cloning Scheme:





^{*} The last codon before the Stop codon of the ORF

ACCN: NM_001135748

ORF Size: 648 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).





Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001135748.3</u>

 RefSeq ORF:
 651 bp

 Locus ID:
 252969

 UniProt ID:
 Q96952

Cytogenetics: 8p23.1

Protein Families: Druggable Genome
Protein Pathways: Base excision repair

MW: 24 kDa


Gene Summary: This gene encodes a member of the Fpg/Nei family of DNA glycosylases. These glycosylases

initiate the first step in base excision repair by cleaving oxidatively damaged bases and introducing a DNA strand break via their abasic site lyase activity. This enzyme is primarily associated with DNA repair during transcription and acts prefentially on cytosine-derived lesions, particularly 5-hydroxyuracil and 5-hydroxycytosine. It contains an N-terminal catalytic domain, a hinge region, and a C-terminal DNA-binding domain with helix-two-turn-helix and zinc finger motifs. This enzyme interacts with the X-ray cross complementing factor 1 scaffold protein as part of a multi-protein DNA repair complex. A pseudogene of this gene has been


identified. [provided by RefSeq, Mar 2017]



Product images:



Circular map for RC226612



Western blot validation of overexpression lysate (Cat# [LY427694]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with RC226612 using transfection reagent MegaTran 2.0 (Cat# [TT210002]).