

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for RC226666L3V

NCOA4 (NM_001145260) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	NCOA4 (NM_001145260) Human Tagged ORF Clone Lentiviral Particle
Symbol:	NCOA4
Synonyms:	ARA70; ELE1; PTC3; RFG
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_001145260
ORF Size:	1950 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC226666).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 001145260.1, NP 001138732.1</u>
RefSeq ORF:	1953 bp
Locus ID:	8031
UniProt ID:	<u>Q13772</u>
Cytogenetics:	10q11.22
Protein Families:	Druggable Genome, Transcription Factors
Protein Pathways:	Pathways in cancer, Thyroid cancer
MW:	73.5 kDa



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Gene Summary:This gene encodes an androgen receptor coactivator. The encoded protein interacts with the
androgen receptor in a ligand-dependent manner to enhance its transcriptional activity.
Chromosomal translocations between this gene and the ret tyrosine kinase gene, also located
on chromosome 10, have been associated with papillary thyroid carcinoma. Alternatively
spliced transcript variants have been described. Pseudogenes are present on chromosomes
4, 5, 10, and 14. [provided by RefSeq, Feb 2009]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US