

Product datasheet for RC226826L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

RBM24 (NM_001143942) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: RBM24 (NM_001143942) Human Tagged ORF Clone Lentiviral Particle

Symbol: RBM24

Synonyms: dJ259A10.1; RNPC6

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_001143942

ORF Size: 708 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC226826).

Sequence:
OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 001143942.1

 RefSeq ORF:
 711 bp

 Locus ID:
 221662

 UniProt ID:
 Q9BX46

 Cytogenetics:
 6p22.3

Protein Families: Druggable Genome

MW: 24.6 kDa





Gene Summary:

Multifunctional RNA-binding protein involved in the regulation of pre-mRNA splicing, mRNA stability and mRNA translation important for cell fate decision and differentiation (PubMed:20977548, PubMed:24375645, PubMed:29358667, PubMed:29104163). Plays a major role in pre-mRNA alternative splicing regulation (PubMed:26990106, PubMed:29104163). Mediates preferentially muscle-specific exon inclusion in numerous mRNAs important for striated cardiac and skeletal muscle cell differentiation (PubMed:29104163). Binds to intronic splicing enhancer (ISE) composed of stretches of GUrich motifs localized in flanking intron of exon that will be included by alternative splicing (By similarity). Involved in embryonic stem cell (ESC) transition to cardiac cell differentiation by promoting pre-mRNA alternative splicing events of several pluripotency and/or differentiation genes (PubMed:26990106). Plays a role in the regulation of mRNA stability (PubMed:20977548, PubMed:24356969, PubMed:24375645, PubMed:29104163). Binds to 3'untranslated region (UTR) AU-rich elements in target transcripts, such as CDKN1A and MYOG, leading to maintain their stabilities (PubMed:20977548, PubMed:24356969). Involved in myogenic differentiation by regulating MYOG levels (PubMed:20977548). Binds to multiple regions in the mRNA 3' UTR of TP63 isoform 2, hence inducing its destabilization (PubMed:24375645). Promotes also the destabilization of the CHRM2 mRNA via its binding to a region in the coding sequence (PubMed:29104163). Plays a role in the regulation of mRNA translation (PubMed:29358667). Mediates repression of p53/TP53 mRNA translation through its binding to U-rich element in the 3' UTR, hence preventing EIF4E from binding to p53/TP53 mRNA and translation initiation (PubMed:29358667). Binds to a huge amount of mRNAs (PubMed:29104163). Required for embryonic heart development, sarcomer and M-band formation in striated muscles (By similarity).[UniProtKB/Swiss-Prot Function]