



Product datasheet for RC226858L3

SPATA2 (NM_001135773) Human Tagged Lenti ORF Clone

Product data:

Product Type:	Expression Plasmids
Tag:	Myc-DDK
Symbol:	SPATA2
Synonyms:	PD1; PPP1R145; tamo
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
E. coli Selection:	Chloramphenicol (34 ug/mL)
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC226858).
Restriction Sites:	SgfI-MluI
Cloning Scheme:	

Cloning sites used for ORF Shuttling:



* The last codon before the Stop codon of the ORF.

ACCN:	NM_001135773
ORF Size:	1560 bp



OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
Note:	Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.
RefSeq:	NM_001135773.1 , NP_001129245.1
RefSeq Size:	4156 bp
RefSeq ORF:	1563 bp
Locus ID:	9825
UniProt ID:	Q9UM82
Cytogenetics:	20q13.13
MW:	58.4 kDa
Gene Summary:	Bridging factor that mediates the recruitment of CYLD to the LUBAC complex, thereby regulating TNF-alpha-induced necroptosis (PubMed:27307491, PubMed:27458237, PubMed:27545878, PubMed:27591049). Acts as a direct binding intermediate that bridges RNF31/HOIP, the catalytic subunit of the LUBAC complex, and the deubiquitinase (CYLD), thereby recruiting CYLD to the TNF-R1 signaling complex (TNF-RSC) (PubMed:27458237, PubMed:27545878, PubMed:27591049). Required to activate the 'Met-1'- (linear) and 'Lys-63'-linked deubiquitinase activities of CYLD (PubMed:27458237, PubMed:27591049). Controls the kinase activity of RIPK1 and TNF-alpha-induced necroptosis by promoting 'Met-1'-linked deubiquitination of RIPK1 by CYLD (By similarity).[UniProtKB/Swiss-Prot Function]