

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for RC227301L3V

SORBS2 (NM_001145673) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	SORBS2 (NM_001145673) Human Tagged ORF Clone Lentiviral Particle
Symbol:	SORBS2
Synonyms:	ARGBP2; PRO0618
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_001145673
ORF Size:	2472 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC227301).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 001145673.1</u>
RefSeq ORF:	2475 bp
Locus ID:	8470
UniProt ID:	<u>094875</u>
Cytogenetics:	4q35.1
MW:	91.1 kDa



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Gene Summary:Arg and c-Abl represent the mammalian members of the Abelson family of non-receptor
protein-tyrosine kinases. They interact with the Arg/Abl binding proteins via the SH3 domains
present in the carboxy end of the latter group of proteins. This gene encodes the sorbin and
SH3 domain containing 2 protein. It has three C-terminal SH3 domains and an N-terminal
sorbin homology (SoHo) domain that interacts with lipid raft proteins. The subcellular
localization of this protein in epithelial and cardiac muscle cells suggests that it functions as
an adapter protein to assemble signaling complexes in stress fibers, and that it is a potential
link between Abl family kinases and the actin cytoskeleton. Alternative splicing results in
multiple transcript variants encoding different isoforms. [provided by RefSeq, Jul 2008]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US