

Product datasheet for RC227442L2V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Bestrophin (BEST1) (NM 001139443) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Bestrophin (BEST1) (NM 001139443) Human Tagged ORF Clone Lentiviral Particle

Symbol: BEST1

Synonyms: ARB; BEST; Best1V1Delta2; BMD; RP50; TU15B; VMD2

Mammalian Cell

Selection:

None

Vector: pLenti-C-mGFP (PS100071)

Tag: mGFP

ACCN: NM_001139443

ORF Size: 1812 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC227442).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 001139443.1, NP 001132915.1

 RefSeq ORF:
 1815 bp

 Locus ID:
 7439

 UniProt ID:
 076090

Cytogenetics: 11q12.3

Protein Families: Druggable Genome, Ion Channels: Other, Transmembrane

MW: 68.9 kDa





Gene Summary:

This gene encodes a member of the bestrophin gene family. This small gene family is characterized by proteins with a highly conserved N-terminus with four to six transmembrane domains. Bestrophins may form chloride ion channels or may regulate voltage-gated L-type calcium-ion channels. Bestrophins are generally believed to form calcium-activated chloride-ion channels in epithelial cells but they have also been shown to be highly permeable to bicarbonate ion transport in retinal tissue. Mutations in this gene are responsible for juvenile-onset vitelliform macular dystrophy (VMD2), also known as Best macular dystrophy, in addition to adult-onset vitelliform macular dystrophy (AVMD) and other retinopathies. Alternative splicing results in multiple variants encoding distinct isoforms. [provided by RefSeq, Nov 2008]