

## Product datasheet for RC227527L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## EXOC7 (NM\_001145299) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

Product Name: EXOC7 (NM 001145299) Human Tagged ORF Clone Lentiviral Particle

Symbol: EXOC7

Synonyms: 2-5-3p; BLOM4; EX070; EX070; EX070p; EXOC1; NEDSEBA; YJL085W

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK

**ACCN:** NM\_001145299

ORF Size: 2121 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC227527).

Sequence:

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 001145299.1

 RefSeq ORF:
 2124 bp

 Locus ID:
 23265

 UniProt ID:
 Q9UPT5

 Cytogenetics:
 17q25.1

**Protein Families:** Druggable Genome

**Protein Pathways:** Insulin signaling pathway

**MW:** 80.5 kDa







## **Gene Summary:**

The protein encoded by this gene is a component of the exocyst complex. The exocyst complex plays a critical role in vesicular trafficking and the secretory pathway by targeting post-Golgi vesicles to the plasma membrane. The encoded protein is required for assembly of the exocyst complex and docking of the complex to the plasma membrane. The encoded protein may also play a role in pre-mRNA splicing through interactions with pre-mRNA-processing factor 19. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene, and a pseudogene of this gene is located on the long arm of chromosome 4. [provided by RefSeq, Nov 2011]