

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for RC227571L3V

DAZAP2 (NM_001136268) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	DAZAP2 (NM_001136268) Human Tagged ORF Clone Lentiviral Particle
Symbol:	DAZAP2
Synonyms:	PRTB
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_001136268
ORF Size:	258 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC227571).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 001136268.1</u>
RefSeq ORF:	261 bp
Locus ID:	9802
UniProt ID:	<u>Q15038</u>
Cytogenetics:	12q13.13
MW:	9 kDa



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Gene Summary:This gene encodes a proline-rich protein which interacts with the deleted in azoospermia
(DAZ) and the deleted in azoospermia-like gene through the DAZ-like repeats. This protein
also interacts with the transforming growth factor-beta signaling molecule SARA (Smad
anchor for receptor activation), eukaryotic initiation factor 4G, and an E3 ubiquitinase that
regulates its stability in splicing factor containing nuclear speckles. The encoded protein may
function in various biological and pathological processes including spermatogenesis, cell
signaling and transcription regulation, formation of stress granules during translation arrest,
RNA splicing, and pathogenesis of multiple myeloma. Multiple transcript variants encoding
different isoforms have been found for this gene. [provided by RefSeq, Oct 2008]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US