

Product datasheet for RC227875L3

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

DC SIGN (CD209) (NM_001144894) Human Tagged Lenti ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: DC SIGN (CD209) (NM_001144894) Human Tagged Lenti ORF Clone

Tag: Myc-DDK
Symbol: DC SIGN

Synonyms: CDSIGN; CLEC4L; DC-SIGN; DC-SIGN1; hDC-SIGN

Mammalian Cell Puromycin

Selection:

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

E. coli Selection: Chloramphenicol (34 ug/mL)

ORF Nucleotide The ORF insert of this clone is exactly the same as(RC227875).

Sequence:

Restriction Sites: Sgfl-Mlul

Cloning Scheme:





^{*} The last codon before the Stop codon of the ORF.

ACCN: NM_001144894

ORF Size: 1080 bp



DC SIGN (CD209) (NM_001144894) Human Tagged Lenti ORF Clone - RC227875L3

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001144894.1, NP 001138366.1</u>

 RefSeq Size:
 4196 bp

 RefSeq ORF:
 1083 bp

 Locus ID:
 30835

 UniProt ID:
 Q9NNX6

 Cytogenetics:
 19p13.2

Protein Families: Druggable Genome

MW: 41 kDa

Gene Summary: This gene encodes a C-type lectin that functions in cell adhesion and pathogen recognition.

impact on public health, including leprosy and tuberculosis mycobacteria, the Ebola, hepatitis C, HIV-1 and Dengue viruses, and the SARS-CoV acute respiratory syndrome coronavirus. The protein is organized into four distinct domains: a C-terminal carbohydrate recognition domain, a flexible tandem-repeat neck domain, a transmembrane region and an N-terminal cytoplasmic domain involved in internalization. This gene is closely related in terms of both sequence and function to a neighboring gene, CLEC4M (Gene ID: 10332), also known as L-SIGN. The two genes differ in viral recognition and expression patterns, with this gene showing high expression on the surface of dendritic cells. Polymorphisms in the neck region are associated with protection from HIV-1 infection, while single nucleotide polymorphisms in the promoter of this gene are associated with differing resistance and susceptibility to and

severity of infectious disease, including rs4804803, which is associated with SARS severity.

This receptor recognizes a wide range of evolutionarily divergent pathogens with a large

[provided by RefSeq, May 2020]