

## Product datasheet for RC227916L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **NEIL2 (NM\_001135747) Human Tagged ORF Clone Lentiviral Particle**

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** NEIL2 (NM\_001135747) Human Tagged ORF Clone Lentiviral Particle

Symbol: NEIL2

Synonyms: NEH2; NEI2

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK

**ACCN:** NM\_001135747

ORF Size: 813 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC227916).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 001135747.1

 RefSeq Size:
 2062 bp

 RefSeq ORF:
 816 bp

 Locus ID:
 252969

 UniProt ID:
 Q969S2

 Cytogenetics:
 8p23.1

Protein Families: Druggable Genome
Protein Pathways: Base excision repair





ORIGENE

**MW:** 29.9 kDa

**Gene Summary:** 

This gene encodes a member of the Fpg/Nei family of DNA glycosylases. These glycosylases initiate the first step in base excision repair by cleaving oxidatively damaged bases and introducing a DNA strand break via their abasic site lyase activity. This enzyme is primarily associated with DNA repair during transcription and acts prefentially on cytosine-derived lesions, particularly 5-hydroxyuracil and 5-hydroxycytosine. It contains an N-terminal catalytic domain, a hinge region, and a C-terminal DNA-binding domain with helix-two-turn-helix and zinc finger motifs. This enzyme interacts with the X-ray cross complementing factor 1 scaffold protein as part of a multi-protein DNA repair complex. A pseudogene of this gene has been identified. [provided by RefSeq, Mar 2017]