

Product datasheet for RC228006L2V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

IL12A (NM_000882) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: IL12A (NM 000882) Human Tagged ORF Clone Lentiviral Particle

Symbol: IL12A

Synonyms: CLMF; IL-12A; NFSK; NKSF1; P35

Mammalian Cell

Selection:

None

Vector: pLenti-C-mGFP (PS100071)

Tag: mGFP

ACCN: NM_000882

ORF Size: 759 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC228006).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 000882.2

 RefSeq Size:
 1450 bp

 RefSeq ORF:
 762 bp

 Locus ID:
 3592

 UniProt ID:
 P29459

Cytogenetics: 3q25.33

Protein Families: Druggable Genome, Secreted Protein



IL12A (NM_000882) Human Tagged ORF Clone Lentiviral Particle - RC228006L2V

Protein Pathways: Allograft rejection, Cytokine-cytokine receptor interaction, Jak-STAT signaling pathway, RIG-I-

like receptor signaling pathway, Toll-like receptor signaling pathway, Type I diabetes mellitus

MW: 28.3 kDa

Gene Summary: This gene encodes a subunit of a cytokine that acts on T and natural killer cells, and has a

broad array of biological activities. The cytokine is a disulfide-linked heterodimer composed of the 35-kD subunit encoded by this gene, and a 40-kD subunit that is a member of the cytokine receptor family. This cytokine is required for the T-cell-independent induction of interferon (IFN)-gamma, and is important for the differentiation of both Th1 and Th2 cells. The responses of lymphocytes to this cytokine are mediated by the activator of transcription protein STAT4. Nitric oxide synthase 2A (NOS2A/NOS2) is found to be required for the

signaling process of this cytokine in innate immunity. [provided by RefSeq, Jul 2008]